Datasheet: LNK162STR

BATCH NUMBER 165496

Name:	STREPTAVIDIN CONJUGATION KIT
Format:	Kit
Product Type:	Conjugation Kit
Quantity:	1 CONJUGATION for 1mg antibody

Applications	This product has been re	anorted to	work in t	he following application	ne This information is			
Αρριτσατιστισ	This product has been reported to work in the following applications. This information is							
	derived from testing within our laboratories, peer-reviewed publications or personal							
	communications from the originators. Please refer to references indicated for further							
	information. For general protocol recommendations, please visit <u>www.bio-</u>							
	rad-antibodies.com/protocols.							
		Yes	No	Not Determined	Suggested Dilution			
	Conjugation	-						
	We recommend that for each conjugation the user determines the best antibody:conjugate							
	ratio.							
	1410.							
Product Information	IVNV Danid Strantovid	in Antiha	dy Caniu	nation Kit@ anablaa	the repid conjugation of			
Product information	LYNX Rapid Streptavid			-				
	a pre-prepared lyophilize	ed mixture	containir	ng Streptavidin label to	o an antibody or protein.			
	Activation of proprietary reagents within the antibody-label solution results in the coupling							
	of Streptavidin to the antibody.							
	•	-						
	LYNX Rapid Streptavidin Antibody Conjugation Kit® can be used to label small quantities							
	I VNX Rapid Streptavidir	a Antihody	Coniuga	tion Kit® can be used	to label small quantities			
		-			•			
	of antibody/protein at ne	-			•			
		-			•			
Reagents in The Kit	of antibody/protein at ne antibody recovery.	ar neutral	pH, allow	ring a high conjugatior	•			
Reagents In The Kit	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop	ar neutral hilized Str	pH, allow	ring a high conjugatior	•			
Reagents In The Kit	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea	ar neutral hilized Str	pH, allow	ring a high conjugatior	•			
Reagents In The Kit	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop	ar neutral hilized Str	pH, allow	ring a high conjugatior	•			
	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea	ar neutral hilized Str agent eagent.	pH, allow	ring a high conjugatior	n efficiency with 100%			
Preparing The	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea The following buffer solu	ar neutral hilized Str agent eagent. itions are i	pH, allow eptavidin recomme	ring a high conjugation mix nded for preparing the	n efficiency with 100%			
Preparing The	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea The following buffer solu 10-50mM amine-free bu	ar neutral hilized Str agent eagent. itions are i	pH, allow reptavidin recomme EPES, M	ring a high conjugation mix nded for preparing the ES, MOPS and phosp	n efficiency with 100% e antibody: ohate) pH range 6.5-8.5,			
Preparing The	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea The following buffer solu	ar neutral hilized Str agent eagent. itions are i	pH, allow reptavidin recomme EPES, M	ring a high conjugation mix nded for preparing the ES, MOPS and phosp	n efficiency with 100% e antibody: ohate) pH range 6.5-8.5,			
Reagents In The Kit Preparing The Antibody	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea The following buffer solu 10-50mM amine-free bu	ar neutral hilized Str agent eagent. ttions are r ffer (e.g H entrations	pH, allow reptavidin recomme EPES, M of Tris bu	nded for preparing the ES, MOPS and phosp	e antibody: bhate) pH range 6.5-8.5, tolerated.			
Preparing The	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea The following buffer solu 10-50mM amine-free bur although moderate conc	ar neutral hilized Str agent eagent. tions are n ffer (e.g H entrations aining nucl	pH, allow reptavidin EPES, M of Tris bu eophilic c	ning a high conjugation mix nded for preparing the ES, MOPS and phosp uffer (<20mM) may be components e.g prima	e antibody: ohate) pH range 6.5-8.5, tolerated. ry amines and thiols			
Preparing The	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea The following buffer solu 10-50mM amine-free bur although moderate conc Do not use buffers conta since they may react wit	ar neutral hilized Str agent eagent. Itions are r ffer (e.g H entrations aining nucl h LYNX ch	pH, allow reptavidin EPES, M of Tris bi eophilic c nemicals.	nded for preparing the ES, MOPS and phosp uffer (<20mM) may be components e.g prima Azide (0.02-0.1%), El	e antibody: bhate) pH range 6.5-8.5, tolerated. ry amines and thiols DTA and common			
Preparing The	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea The following buffer solu 10-50mM amine-free bur although moderate conc Do not use buffers conta since they may react with non-buffering salts and s	ar neutral chilized Str agent eagent. tions are r ffer (e.g H entrations aining nucl h LYNX ch sugars hav	pH, allow reptavidin EPES, M of Tris bu eophilic c nemicals. /e little or	nded for preparing the ES, MOPS and phosp uffer (<20mM) may be components e.g prima Azide (0.02-0.1%), El no effect on conjugati	e antibody: ohate) pH range 6.5-8.5, tolerated. ry amines and thiols DTA and common ion efficiency.			
Preparing The	of antibody/protein at ne antibody recovery. 1 Vial of 1mg LYNX lyop 1 Vial LYNX Modifier rea 1 Vial LYNX Quencher rea The following buffer solu 10-50mM amine-free bur although moderate conc Do not use buffers conta since they may react wit	ar neutral hilized Str agent eagent. Itions are r ffer (e.g H entrations aining nucl h LYNX ch sugars hav	pH, allow reptavidin EPES, M of Tris bu eophilic c nemicals. /e little or pdy be us	nded for preparing the ES, MOPS and phosp uffer (<20mM) may be components e.g prima Azide (0.02-0.1%), EI no effect on conjugati ed in each labelling re	e antibody: ohate) pH range 6.5-8.5, tolerated. ry amines and thiols DTA and common ion efficiency. eaction. For optimal			

	1ml and a maximum antibody amount of 1mg. Antibody at a concentration of greater than 1mg/ml requires dilution. Antibody below 1mg/ml can still be used as long as the maximum volume is not exceeded. Using less than the recommended amount of antibody may result in unbound label, but this will be removed during subsequent application wash steps. Antibody below 0.5mg/ml should be concentrated before use with the kit.
Instructions For Use	 1.To the antibody sample add 1µl of the Modifier reagent for every 10µl of antibody and mix gently. 2.Pipette the mixed antibody-modifier sample directly onto the LYNX lyophilized mix and gently pipette up and down twice to resuspend. 3.Replace cap onto vial and incubate at room temperature (20-25°C) for 3 hours or overnight.if preferred.
	4.After incubation, add 1µl of Quencher reagent for every 10µl of antibody used. Leave to stand for 30 minutes before use.
References	1. Welden, M. <i>et al.</i> (2022) Towards Multi-Analyte Detection with Field-Effect Capacitors Modified with <i>Tobacco Mosaic Virus</i> Bioparticles as Enzyme Nanocarriers. <u>Biosensors</u> (Basel). 12 (1): 43.
Storage	This kit contains lyophilized hygroscopic components that are moisture-sensitive. This kit is shipped under ambient conditions with silica packets to avoid exposure to moisture. On receipt, Bio-Rad recommend that the kit is stored at -20°C and protected from moisture. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing. Before opening, allow the components to reach room temperature to minimize condensation.
Guarantee	12 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10536 #10546 #10548 available at: <u>https://www.bio-rad-antibodies.com/SDS/LNK162STR</u> Lyophilized Streptavidin Mix (10536) Modifier Reagent (10546) Quencher Reagent (10548)
Licensed Use	Bio-Rad LYNX conjugation kits are offered for research purposes alone, and are not intended for human, therapeutic or diagnostic use. The purchase of this conjugation kit conveys to the buyer (whether the buyer is a not-for-profit, academic or for-profit entity) the non-transferable right to use the amount of product purchased and the components of the product for in-house research. The buyer shall not sell or otherwise transfer this product, its components, or materials prepared therefrom to any third party. The buyer shall not use this product or its components for commercial purposes. For the avoidance of doubt, ?commercial purposes? means any activity by a party for consideration and includes, without limitation, use of the product or its components (I) in the manufacturing of conjugated materials (e.g. labeled antibodies), (ii) to provide a service, information or data, (iii) for therapeutic, diagnostic or prophylactic purposes, or (iv) for repackaging/resale, whether or not such product or its components are resold for use in research. The use of this product by the buyer constitutes agreement with the terms of this limited use label license for LYNX products.

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-ra	Worldwide ad.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rac	Europe d.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot specific datasheet	for this produc	ct, please use our online sea 'M404970:220914'	arch tool at: b	io-rad-antibodies.com/datasheets

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint