## Datasheet: LNK174PETR BATCH NUMBER 166306

LYNX RAPID	RPF-Texas R					
			BODT CONJUGATIO			
RPE-Texas Re						
Kit						
Conjugation K	lit					
3 CONJUGAT	IONS for 10µ	g antibody	/			
This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.						
Conjugation	•	NO	Not Determined	Suggested Dilution		
We recommend that ratio	for each conj	ugation th	e user determines the	e best antibody:conjugate		
Product InformationLYNX Rapid RPE-Texas Red® Antibody Conjugation Kit® enables the rapid conjugation of a pre-prepared lyophilized mixture containing R-Phycoerythrin (R Red® label to an antibody or protein. Activation of proprietary reagents within th antibody-label solution results in directional covalent bonding of RPE-Texas Red antibody.						
The LYNX Rapid Conjugation kit® can be used to label small quantities of antibody/protein at near neutral pH, allowing a high conjugation efficiency with 100% antibody recovery.						
1 Vial LYNX Modifier	reagent.	RPE-Texa	s Red mix.			
The following buffer solutions are recommended for preparing the antibody:						
10-50mM amine-free buffer (e.g HEPES, MES, MOPS and phosphate) pH range 6.5-8.5, although moderate concentrations of Tris buffer (<20mM) may be tolerated.						
If possible, avoid buffers containing nucleophilic components such as primary amines and thiols (e.g. thiomersal/thimerosal) since they may react with LYNX						
	Kit   Conjugation K   3 CONJUGAT   This product has been derived from testing communications from information. For generad-antibodies.com/p   Conjugation   Ke recommend that ratio   LYNX Rapid RPE-Teconjugation of a pre-Red® label to an antibody-label solution antibody.   The LYNX Rapid Co antibody/protein at n antibody recovery.   3 Vials of 10µg LYNX 1 Vial LYNX Modifier 1 Vial LYNX Quench   The following buffer   10-50mM amine-free although moderate combined for the combined for t	Kit   Conjugation Kit   3 CONJUGATIONS for 10µg   This product has been reported to derived from testing within our labor communications from the originator information. For general protocol report rad-antibodies.com/protocols.   Yes   Conjugation   We recommend that for each conju- ratio   LYNX Rapid RPE-Texas Red® Age conjugation of a pre-prepared lyop Red® label to an antibody or prote- antibody-label solution results in d antibody.   The LYNX Rapid Conjugation kit® antibody/protein at near neutral ph- antibody recovery.   3 Vials of 10µg LYNX lyophilized F 1 Vial LYNX Modifier reagent.   1 Vial LYNX Quencher reagent.   1 Vial LYNX Quencher reagent.   10-50mM amine-free buffer (e.g H although moderate concentrations   If possible, avoid buffers contail	Kit   Conjugation Kit   3 CONJUGATIONS for 10µg antibody   This product has been reported to work in the derived from testing within our laboratories, communications from the originators. Please information. For general protocol recomment rad-antibodies.com/protocols.   Yes No   Conjugation •   We recommend that for each conjugation the ratio •   LYNX Rapid RPE-Texas Red® Antibody C conjugation of a pre-prepared lyophilized mix Red® label to an antibody or protein. Activa antibody-label solution results in directional antibody.   The LYNX Rapid Conjugation kit® can be us antibody/protein at near neutral pH, allowing antibody recovery.   3 Vials of 10µg LYNX lyophilized RPE-Texas: 1 Vial LYNX Modifier reagent.   1 Vial LYNX Quencher reagent.   1 Vial LYNX Quencher reagent.   10-50mM amine-free buffer (e.g HEPES, Mi although moderate concentrations of Tris buffer solutions and the solution results in a solution soluti	Conjugation Kit   3 CONJUGATIONS for 10µg antibody   This product has been reported to work in the following application derived from testing within our laboratories, peer-reviewed publica communications from the originators. Please refer to references in information. For general protocol recommendations, please visit y rad-antibodies.com/protocols.   Yes No   Not Determined   Conjugation •   We recommend that for each conjugation the user determines the ratio   LYNX Rapid RPE-Texas Red® Antibody Conjugation Kit® ena conjugation of a pre-prepared lyophilized mixture containing R-PhRed® label to an antibody or protein. Activation of proprietary rea antibody-label solution results in directional covalent bonding of R antibody.   The LYNX Rapid Conjugation kit® can be used to label small qua antibody recovery.   3 Vials of 10µg LYNX lyophilized RPE-Texas Red mix.   1 Vial LYNX Quencher reagent.   1 Vial LYNX Quencher reagent.   10-50mM amine-free buffer (e.g HEPES, MES, MOPS and phosp although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of Tris buffer (<20mM) may be although moderate concentrations of T		

	<b>chemicals</b> . Azide (0.02-0.1%), EDTA, up to 50% Glycerol and common non-buffering salts and sugars have little or no effect on conjugation efficiency.				
	For optimal results the antibody should be at a concentration of 1mg/ml, with a maximum volume of 10ul and a maximum antibody amount of 10µg. Antibody at a concentration of greater than 1mg/ml requires dilution. Antibody below 1mg/ml can still be used as long as the maximum volume is not exceeded. Using less than the recommended amount of antibody may result in unbound label, but this will be removed during subsequent application wash steps. Antibody below 0.5mg/ml should be concentrated before use with the kit.				
Instructions For Use	1. To the antibody sample add $1\mu$ I of the Modifier reagent for every $10\mu$ I of antibody and mix gently.				
	2. Pipette the mixed antibody-modifier sample directly onto the LYNX lyophilized mix and gently pipette up and down twice to resuspend.				
	3. Replace cap onto vial and incubate in the dark at room temperature (20-25°C) for 3 hours, or overnight if preferred.				
	4. After incubation, add 1 $\mu$ l of Quencher reagent for every $\mu$ l of antibody used. Leave to stand for 30 minutes before use.				
Storage	This kit contains lyophilized hygroscopic components that are moisture-sensitive. This kit is shipped under ambient conditions with silica packets to avoid exposure to moisture. On receipt, Bio-Rad recommend that the kit is stored at -20°C and protected from moisture. Storage in frost-free freezers is not recommended.This product should be stored undiluted. Avoid repeated freezing and thawing. Before opening, allow the components to reach room temperature to minimize condensation.				
Guarantee	12 months from date of despatch				
Acknowledgements	Texas Red is a trademark of Molecular Probes, Inc.				
Health And Safety Information	Material Safety Datasheet documentation #10537 #10546 #10549 available at: <u>https://www.bio-rad-antibodies.com/SDS/LNK174PETR</u> Lyophilized RPE-Texas Red Mix (10537) Modifier Reagent (10546) Quencher Reagent (10549)				
Licensed Use	Bio-Rad LYNX Rapid Conjugation kits are offered for research purposes alone, and are not intended for human, therapeutic or diagnostic use. The purchase of this conjugation kit conveys to the buyer (whether the buyer is a not-for-profit, academic or for-profit entity) the non-transferable right to use the amount of product purchased and the components of the product for in-house research. The buyer shall not sell or otherwise transfer this product, its components, or materials prepared therefrom to any third party. The buyer shall not use this product or its components for commercial purposes. For the avoidance				

includes, without limitation, use of the product or its components (I) in the manufacturing of conjugated materials (e.g. labeled antibodies), (ii) to provide a service, information or data, (iii) for therapeutic, diagnostic or prophylactic purposes, or (iv) for repackaging/resale, whether or not such product or its components are resold for use in research. The use of this product by the buyer constitutes agreement with the terms of this limited use label license for LYNX products.

Regulato	<b>For res</b>				
North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
America	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot specific datash	eet for this produ	uct, please use our online 'M407747:221009'	search tool at	: bio-rad-antibodies.com/datasheets

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint