Datasheet: MCA02R BATCH NUMBER 147840

Description:	MOUSE ANTI MOUSE CD90
Specificity:	CD90
Other names:	THY1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	F7D5
Isotype:	IgM
Quantity:	0.25 mg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

Yes	No	Not Determined	Suggested Dilution
-			1/10 - 1/25
		•	
		•	
		•	
		•	
		•	
-			
	•	•	•

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	IgM fraction - liquid
Preparation	IgM fraction prepared by ammonium sulphate precipitation
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Approx. Protein Concentrations	IgM concentration 1.0 mg/ml
External Database Links	UniProt: P01831 Related reagents Entrez Gene: 21838 Thy1 Related reagents
Synonyms	Thy-1
RRID	AB_323481
Fusion Partners	Spleen cells from immunised AKR mice were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	Mouse anti Mouse CD90 antibody, clone F7D5 recognizes the mouse Thy1.2 alloantigen, also known as CD90.2, which is expressed by thymocytes and peripheral T lymphocytes. Clone F7D5 reacts with Thy1.2 mice such as CBA and BALB/C, but not with Thy1.1 mice eg. AKR and FUB. The antibody is particularly useful for removal of T lymphocytes from cell populations by complement mediated cytotoxicity (Lake <i>et al.</i> 1979).
	Mouse anti Mouse CD90 antibody, clone F7D5 is routinely tested in flow cytometry using mouse thymocytes.
References	 Lake, P. <i>et al.</i> (1979) Production and characterization of cytotoxic Thy-1 antibody- secreting hybrid cell lines. Detection of T cell subsets. <u>Eur J Immunol. 9 (11): 875-86.</u> Yoshida, K. <i>et al.</i> (2002) Evidence for shared recognition of a peptide ligand by a diverse panel of non-obese diabetic mice-derived, islet-specific, diabetogenic T cell clones. <u>Int Immunol. 14 (12): 1439-47.</u> DeVries-vanDerZwan, A. <i>et al.</i> (1997) Specific tolerance induction and transplantation: a single-day protocol. <u>Blood. 89 (7): 2596-601.</u> Winzeler, A.M. <i>et al.</i> (2011) The lipid sulfatide is a novel myelin-associated inhibitor of CNS axon outgrowth. <u>J Neurosci. 31: 6481-92.</u> Unterlauft, J.D. <i>et al.</i> (2014) Enhanced survival of retinal ganglion cells is mediated by Müller glial cell-derived PEDF. <u>Exp Eye Res. 127: 206-14.</u> Brown, R.L. <i>et al.</i> (2015) TRPM3 Expression in Mouse Retina. <u>PLoS One. 10: e0117615.</u> Hanafusa, T. <i>et al.</i> (1988) Induction of insulitis by adoptive transfer with L3T4+Lyt2- T-lymphocytes in T-lymphocyte-depleted NOD mice. <u>Diabetes. 37: 204-8.</u> Billiau, A.D. <i>et al.</i> (2003) Transient expansion of Mac1+Ly6-G+Ly6-C+ early myeloid cells with suppressor activity in spleens of murine radiation marrow chimeras: possible implications for the graft-versus-host and graft-versus-leukemia reactivity of donor lymphocyte infusions. <u>Blood. 102: 740-8.</u> Raeber, A.J. <i>et al.</i> (1999) PrP-dependent association of prions with splenic but not

circulating lymphocytes of scrapie-infected mice. EMBO J. 18: 2702-6.

10. Wang, X. *et al.* (2001) Functional soluble CD100/Sema4D released from activated lymphocytes: possible role in normal and pathologic immune responses. <u>Blood. 97 (11):</u> <u>3498-504.</u>

11. Billiau, A.D. *et al.* (2002) Crucial role of timing of donor lymphocyte infusion in generating dissociated graft-versus-host and graft-versus-leukemia responses in mice receiving allogeneic bone marrow transplants. <u>Blood. 100 (5): 1894-902.</u>

12. Ishikawa, N. *et al.* (1998) Early cytokine responses during intestinal parasitic infections. <u>Immunology. 93 (2): 257-63.</u>

13. Ishigaki, H. *et al.* (2006) Preparation and functional analysis of tumor-infiltrating stroma cells using bone marrow chimera mice. <u>Microbiol Immunol. 50 (8): 655-62.</u>

14. Logan, G.J. *et al.* (2004) CD4 expression on EL4 cells as an epiphenomenon of retroviral transduction and selection. <u>Immunol Cell Biol. 82 (2): 132-5.</u>

15. Gobin, V. *et al.* (2013) Fluoxetine reduces murine graft-versus-host disease by induction of T cell immunosuppression. <u>J Neuroimmune Pharmacol. 8 (4): 934-43.</u>

van Pel, M. *et al.* (2003) Towards a myeloablative regimen with clinical potential: I.
 Treosulfan conditioning and bone marrow transplantation allow induction of donor-specific tolerance for skin grafts across full MHC barriers. <u>Bone Marrow Transplant. 32 (1): 15-22.</u>
 Oosterwegel, M.A. *et al.* (1999) The role of CTLA-4 in regulating Th2 differentiation. <u>J</u>
 Immunol. 163 (5): 2634-9.

18. Vadivelu, S. *et al.* (2015) NG2+ Progenitors Derived From Embryonic Stem Cells Penetrate Glial Scar and Promote Axonal Outgrowth Into White Matter After Spinal Cord Injury. <u>Stem Cells Transl Med. pii: sctm.2014-0107.</u>

19. Wang, Y.L. *et al.* (2015) Electrospun and woven silk fibroin/poly(lactic-co-glycolic acid) nerve guidance conduits for repairing peripheral nerve injury. <u>Neural Regen Res. 10 (10):</u> 1635-42.

20. Liu, X. *et al.* (2017) Thy-1 interaction with Fas in lipid rafts regulates fibroblast apoptosis and lung injury resolution. <u>Lab Invest. (3): 256-67.</u>

21. Naaldijk, Y. *et al.* (2016) Effect of systemic transplantation of bone marrow-derived mesenchymal stem cells on neuropathology markers in APP/PS1 Alzheimer mice. Neuropathol Appl Neurobiol. Feb 26. [Epub ahead of print]

22. Bernard-Marissal, N. *et al.* (2015) Dysfunction in endoplasmic reticulum-mitochondria crosstalk underlies SIGMAR1 loss of function mediated motor neuron degeneration. <u>Brain.</u> <u>138 (Pt 4): 875-90.</u>

23. Takahama, S. *et al.* (2017) Retinal Astrocytes and GABAergic Wide-Field Amacrine Cells Express PDGFRα: Connection to Retinal Ganglion Cell Neuroprotection by PDGF-AA. <u>Invest Ophthalmol Vis Sci. 58 (11): 4703-11.</u>

24. Zhu, B. *et al.* (2019) GAIN domain-mediated cleavage is required for activation of G protein-coupled receptor 56 (GPR56) by its natural ligands and a small-molecule agonist. J Biol Chem. pii: jbc.RA119.008234. Oct 18 [Epub ahead of print].

25. Bürger, S. *et al.* (2020) Pigment Epithelium-Derived Factor (PEDF) Receptors Are Involved in Survival of Retinal Neurons. Int J Mol Sci. 22 (1): 369.

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

	Storage in frost free fr as this may denature recommend microcen	eated freezing and thawing in a precipitate we				
Guarantee	12 months from date of despatch					
Health And Safety Information	Material Safety Datas <u>https://www.bio-rad-ar</u> 10040	at:				
Regulatory	For research purpose					
Related Produce Recommended Second	cts econdary Antibodies					
Goat Anti Mouse Igl Goat Anti Mouse Ig(и (STAR138) <u>All</u> G IgA IgM (STAR87) <u>HF</u>	<u>k. Phos.</u> RP				
North & South Tel: +1 800 263 America Fax: +1 919 87 Email: antibodi		Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad	Europe I.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com		

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M371159:200529'

Printed on 02 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint