

Datasheet: MCA02R

BATCH NUMBER 174547

Description:	MOUSE ANTI MOUSE CD90
Specificity:	CD90
Other names:	THY1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	F7D5
Isotype:	IgM
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/25
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	
Cytotoxic Assays	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	IgM fraction - liquid.
Preparation	IgM fraction prepared by ammonium sulphate precipitation from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline.
Preservative Stabilisers	0.09% sodium azide.

Approx. Protein Concentrations	IgM concentration 1.0 mg/ml.
External Database Links	UniProt: P01831 Related reagents Entrez Gene: 21838 Thy1 Related reagents
Synonyms	Thy-1
RRID	AB_323481
Fusion Partners	Spleen cells from immunized AKR mice were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	Mouse anti Mouse CD90 antibody, clone F7D5 recognizes the mouse Thy1.2 alloantigen, also known as CD90.2, which is expressed by thymocytes and peripheral T lymphocytes. Clone F7D5 reacts with Thy1.2 mice such as CBA and BALB/C, but not with Thy1.1 mice eg. AKR and FUB. The antibody is particularly useful for removal of T lymphocytes from cell populations by complement mediated cytotoxicity (Lake et al. 1979). Mouse anti Mouse CD90 antibody, clone F7D5 is routinely tested in flow cytometry using mouse thymocytes.
References	1. Lake, P. <i>et al.</i> (1979) Production and characterization of cytotoxic Thy-1 antibody-secreting hybrid cell lines. Detection of T cell subsets. Eur J Immunol. 9 (11): 875-86. 2. Hanafusa, T. <i>et al.</i> (1988) Induction of insulinitis by adoptive transfer with L3T4+Lyt2-T-lymphocytes in T-lymphocyte-depleted NOD mice. Diabetes. 37: 204-8. 3. DeVries-vanDerZwan, A. <i>et al.</i> (1997) Specific tolerance induction and transplantation: a single-day protocol. Blood. 89 (7): 2596-601. 4. Ishikawa, N. <i>et al.</i> (1998) Early cytokine responses during intestinal parasitic infections. Immunology. 93 (2): 257-63. 5. Oosterwegel, M.A. <i>et al.</i> (1999) The role of CTLA-4 in regulating Th2 differentiation. J Immunol. 163 (5): 2634-9. 6. Raeber, A.J. <i>et al.</i> (1999) PrP-dependent association of prions with splenic but not circulating lymphocytes of scrapie-infected mice. EMBO J. 18: 2702-6. 7. Wang, X. <i>et al.</i> (2001) Functional soluble CD100/Sema4D released from activated lymphocytes: possible role in normal and pathologic immune responses. Blood. 97 (11): 3498-504. 8. Yoshida, K. <i>et al.</i> (2002) Evidence for shared recognition of a peptide ligand by a diverse panel of non-obese diabetic mice-derived, islet-specific, diabetogenic T cell clones. Int Immunol. 14 (12): 1439-47. 9. Billiau, A.D. <i>et al.</i> (2002) Crucial role of timing of donor lymphocyte infusion in generating dissociated graft-versus-host and graft-versus-leukemia responses in mice

- receiving allogeneic bone marrow transplants. [Blood. 100 \(5\): 1894-902.](#)
10. van Pel, M. *et al.* (2003) Towards a myeloablative regimen with clinical potential: I. Treosulfan conditioning and bone marrow transplantation allow induction of donor-specific tolerance for skin grafts across full MHC barriers. [Bone Marrow Transplant. 32 \(1\): 15-22.](#)
 11. Billiau, A.D. *et al.* (2003) Transient expansion of Mac1+Ly6-G+Ly6-C+ early myeloid cells with suppressor activity in spleens of murine radiation marrow chimeras: possible implications for the graft-versus-host and graft-versus-leukemia reactivity of donor lymphocyte infusions. [Blood. 102: 740-8.](#)
 12. Ishigaki, H. *et al.* (2006) Preparation and functional analysis of tumor-infiltrating stroma cells using bone marrow chimera mice. [Microbiol Immunol. 50 \(8\): 655-62.](#)
 13. Winzeler, A.M. *et al.* (2011) The lipid sulfatide is a novel myelin-associated inhibitor of CNS axon outgrowth. [J Neurosci. 31: 6481-92.](#)
 14. Gobin, V. *et al.* (2013) Fluoxetine reduces murine graft-versus-host disease by induction of T cell immunosuppression. [J Neuroimmune Pharmacol. 8 \(4\): 934-43.](#)
 15. Unterlauff, J.D. *et al.* (2014) Enhanced survival of retinal ganglion cells is mediated by Müller glial cell-derived PEDF. [Exp Eye Res. 127: 206-14.](#)
 16. Vadivelu, S. *et al.* (2015) NG2+ Progenitors Derived From Embryonic Stem Cells Penetrate Glial Scar and Promote Axonal Outgrowth Into White Matter After Spinal Cord Injury. [Stem Cells Transl Med. pii: sctm.2014-0107.](#)
 17. Wang, Y.L. *et al.* (2015) Electrospun and woven silk fibroin/poly(lactic-co-glycolic acid) nerve guidance conduits for repairing peripheral nerve injury. [Neural Regen Res. 10 \(10\): 1635-42.](#)
 18. Bernard-Marissal, N. *et al.* (2015) Dysfunction in endoplasmic reticulum-mitochondria crosstalk underlies SIGMAR1 loss of function mediated motor neuron degeneration. [Brain. 138 \(Pt 4\): 875-90.](#)
 19. Brown, R.L. *et al.* (2015) TRPM3 Expression in Mouse Retina. [PLoS One. 10: e0117615.](#)
 20. Liu, X. *et al.* (2017) Thy-1 interaction with Fas in lipid rafts regulates fibroblast apoptosis and lung injury resolution. [Lab Invest. \(3\): 256-67.](#)
 21. Naaldijk, Y. *et al.* (2017) Effect of systemic transplantation of bone marrow-derived mesenchymal stem cells on neuropathology markers in APP/PS1 Alzheimer mice. [Neuropathol Appl Neurobiol. 43 \(4\): 299-314.](#)
 22. Takahama, S. *et al.* (2017) Retinal Astrocytes and GABAergic Wide-Field Amacrine Cells Express PDGFR α : Connection to Retinal Ganglion Cell Neuroprotection by PDGF-AA. [Invest Ophthalmol Vis Sci. 58 \(11\): 4703-11.](#)
 23. Zhu, B. *et al.* (2019) GAIN domain-mediated cleavage is required for activation of G protein-coupled receptor 56 (GPR56) by its natural ligands and a small-molecule agonist. [J Biol Chem. pii: jbc.RA119.008234. Oct 18 \[Epub ahead of print\].](#)
 24. Bürger, S. *et al.* (2020) Pigment Epithelium-Derived Factor (PEDF) Receptors Are Involved in Survival of Retinal Neurons. [Int J Mol Sci. 22 \(1\): 369.](#)
 25. Qiu, A.W. *et al.* (2021) IL-17A injury to retinal ganglion cells is mediated by retinal Müller cells in diabetic retinopathy. [Cell Death Dis. 12 \(11\): 1057.](#)
 26. Xing, J. *et al.* (2023) Post-injury born oligodendrocytes incorporate into the glial scar and contribute to the inhibition of axon regeneration. [Development. 150 \(8\): dev201311.](#)
 27. Ma, R. *et al.* (2023) RGC-Net: An Automatic Reconstruction and Quantification Algorithm for Retinal Ganglion Cells Based on Deep Learning. [Transl Vis Sci Technol. 12 \(5\): 7.](#)

28. Xing, J. *et al.* (2023) Experimental upregulation of developmentally downregulated ribosomal protein large subunits 7 and 7A promotes axon regeneration after injury *in vivo*. [Exp Neurol. 368: 114510.](#)
29. Llorente, I.L. *et al.* (2021) Patient-derived glial enriched progenitors repair functional deficits due to white matter stroke and vascular dementia in rodents. [Sci Transl Med. 13 \(590\) \[Epub ahead of print\].](#)
30. Lukomska, A. *et al.* (2024) Upregulation of developmentally-downregulated miR-1247-5p promotes neuroprotection and axon regeneration *in vivo*. [Neurosci Lett. 823: 137662.](#)
31. Sun, S. *et al.* (2020) Opa1 Deficiency Leads to Diminished Mitochondrial Bioenergetics With Compensatory Increased Mitochondrial Motility. [Invest Ophthalmol Vis Sci. 61 \(6\): 42.](#)
32. Zwanzig, A. *et al.* (2021) Neuroprotective effects of glial mediators in interactions between retinal neurons and Müller cells. [Exp Eye Res. 209: 108689.](#)
33. Qiu, A.W. *et al.* (2021) IL-17A injury to retinal ganglion cells is mediated by retinal Müller cells in diabetic retinopathy. [Cell Death Dis. 12 \(11\): 1057.](#)
34. Lukomska, A. *et al.* (2024) Augmenting fibronectin levels in injured adult CNS promotes axon regeneration *in vivo*. [Exp Neurol. 379: 114877.](#)
35. Qiu, A.W. *et al.* (2025) Retinal Müller Cell-Released Exosomal MiR-92a-3p Delivers Interleukin-17A Signal by Targeting Notch-1 to Promote Diabetic Retinopathy. [Invest Ophthalmol Vis Sci. 66 \(1\): 1.](#)

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA02R
Regulatory	For research purposes only.

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgM (STAR138...) [Alk. Phos.](#)
 Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
 'M406848:221002'

Printed on 27 Aug 2026

