

Datasheet: MCA04GT

BATCH NUMBER 160810

Description:	MOUSE ANTI RAT CD90
Specificity:	CD90
Other names:	THY1, THY1.1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	T11D7e
Isotype:	IgM
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/5
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Cytotoxic Assays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat
Species Cross Reactivity	<p>Reacts with: Mouse</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Ig fraction - liquid

Preparation	Ig fraction prepared by ammonium sulphate precipitation from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgM concentration 1 mg/ml
Immunogen	Xenogeneic rat thymocytes.
External Database Links	<p>UniProt: P01830 Related reagents</p> <p>Entrez Gene: 24832 Thy1 Related reagents</p>
Synonyms	Thy-1
RRID	AB_2201293
Fusion Partners	Spleen cells from immunised CBA mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	<p>Mouse anti Rat CD90 antibody, clone T11D7e recognizes mouse and rat CD90 also known as Thy1.1. Thy1.1 is a monomorphic determinant in rat but polymorphic in mice. Clone T11D7e reacts with Thy1.1 mice e.g. AKR and FVB mice, but not Thy1.2 mice such as CBA and BALB/c.</p> <p>Clone T11D7e is useful for removal of T lymphocytes from cell populations by complement mediated cytotoxicity (Lake et al. 1979).</p> <p>Mouse anti Rat CD90 antibody, clone T11D7e is routinely tested in flow cytometry using mouse thymocytes.</p>
References	<ol style="list-style-type: none"> 1. Lake, P. <i>et al.</i> (1979) Production and characterization of cytotoxic Thy-1 antibody-secreting hybrid cell lines. Detection of T cell subsets. Eur J Immunol. 9 (11): 875-86. 2. Yasumizu, R. <i>et al.</i> (1988) Development of donor-derived thymic lymphomas after allogeneic bone marrow transplantation in AKR/J mice. J Immunol. 141 (6): 2181-6. 3. Fabian, R.H. (1990) Uptake of antineuronal IgM by CNS neurons: comparison with antineuronal IgG. Neurology. 40 (3 Pt 1): 419-22. 4. Ogasawara, K. <i>et al.</i> (1990) Ia restriction specificity of KLH-specific T cells from allogeneic bone marrow chimeras is influenced by histocompatibility at the H-2 and minor histocompatibility loci. Microbiol Immunol. 34 (12): 1025-39. 5. Mahanthappa, N.K. <i>et al.</i> (1996) Glial growth factor 2, a soluble neuregulin, directly increases Schwann cell motility and indirectly promotes neurite outgrowth. J Neurosci. 16: 4673-83.

6. Simitzi, C. *et al.* (2015) Data in support on the shape of Schwann cells and sympathetic neurons onto microconically structured silicon surfaces. [Data Brief. 4: 636-40.](#)
7. Pasten, C. *et al.* (2015) ApoER2 and Reelin are expressed in regenerating peripheral nerve and regulate Schwann cell migration by activating the Rac1 GEF protein, Tiam1. [Mol Cell Neurosci. 69: 1-11.](#)
8. Deliyanti, D. & Wilkinson-Berka, J.L. (2015) Inhibition of NOX1/4 with GKT137831: a potential novel treatment to attenuate neuroglial cell inflammation in the retina. [J Neuroinflammation. 12: 136.](#)
9. Heinen, A. *et al.* (2015) Fingolimod induces the transition to a nerve regeneration promoting Schwann cell phenotype. [Exp Neurol. 271: 25-35.](#)
10. Dietrich, N. *et al.* (2016) The DPP4 Inhibitor Linagliptin Protects from Experimental Diabetic Retinopathy. [PLoS One. 11 \(12\): e0167853.](#)
11. Ee, X. *et al.* (2017) Transgenic SCs expressing GDNF-IRES-DsRed impair nerve regeneration within acellular nerve allografts. [Biotechnol Bioeng. 114 \(9\): 2121-30.](#)
12. Wang, J. *et al.* (2020) CTCF-mediated chromatin looping in EGR2 regulation and SUZ12 recruitment critical for peripheral myelination and repair. [Nat Commun. 11 \(1\): 4133.](#)
13. Warren, P.M. *et al.* (2020) Secretion of a mammalian chondroitinase ABC aids glial integration at PNS/CNS boundaries. [Sci Rep. 10 \(1\): 11262.](#)
14. Piovesana, R. *et al.* (2020) Muscarinic receptors modulate Nerve Growth Factor production in rat Schwann-like adipose-derived stem cells and in Schwann cells. [Sci Rep. 10 \(1\): 7159.](#)
15. Castelnovo, L.F. *et al.* (2020) Membrane Progesterone Receptors (mPRs/PAQRs) Differently Regulate Migration, Proliferation, and Differentiation in Rat Schwann Cells. [J Mol Neurosci. 70 \(3\): 433-48.](#)
16. Ma, S. *et al.* (2021) Effect of papaverine on axonal outgrowth of primary retinal ganglion cells of Sprague Dawley rats. [Exp Eye Res. 212: 108797.](#)
17. Salazar Intriago, M.S. *et al.* (2021) The Mechanisms Mediated by $\alpha 7$ Acetylcholine Nicotinic Receptors May Contribute to Peripheral Nerve Regeneration. [Molecules.26\(24\):7668.](#)
18. Uezumi, A. *et al.* (2021) Mesenchymal Bmp3b expression maintains skeletal muscle integrity and decreases in age-related sarcopenia. [J Clin Invest. 131 \(1\): e139617.](#)
19. De Vitis, E. *et al.* (2021) A microfabricated multi-compartment device for neuron and Schwann cell differentiation. [Sci Rep. 11 \(1\): 7019.](#)
20. Botticelli, E. *et al.* (2022) Analysis of Signal Transduction Pathways Downstream M2 Receptor Activation: Effects on Schwann Cell Migration and Morphology [Life. 12 \(2\): 211.](#)
21. Piovesana, R. *et al.* (2022) Notch Signal Mediates the Cross-Interaction between M2 Muscarinic Acetylcholine Receptor and Neuregulin/ErbB Pathway: Effects on Schwann Cell Proliferation. [Biomolecules. 12 \(2\): 239.](#)
22. Huang, J.H. *et al.* (2022) Schwann cells-derived exosomes promote functional recovery after spinal cord injury by promoting angiogenesis. [Front Cell Neurosci. 16: 1077071.](#)

Storage

Store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA04GT 10040
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgM (STAR138...) [Alk. Phos.](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M371163:200529'

Printed on 18 Jan 2024