

Datasheet: MCA1014FB BATCH NUMBER 150155

MOUSE ANTI MOUSE CD44:FITC
CD44
H-CAM, PGP-1
FITC
Monoclonal Antibody
5035-41.1D
lgG1

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry	-		1/10 - 1/50		
	Where this antibody ha necessarily exclude its a guide only. It is recon	use in such proced	ures. Suggested workir	ng dilutions are given as		
	system using appropria	te negative/positive	e controls.			
Target Species	Mouse					
Product Form	Purified IgG conjugated	to Fluorescein Isc	thiocyanate Isomer 1 (F	FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm) Emission Max (nm)			
	FITC	490	525			
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered sal	ine				
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum A	lbumin				
Approx. Protein	IgG concentration 0.5 r	ng/ml				

Concentrations

Spleen cells from B6PL-Ly-2a / Ly-3a B6-Ly-1a mice.
UniProt: <u>P15379</u> <u>Related reagents</u>
Entrez Gene: <u>12505</u> Cd44 <u>Related reagents</u>
Ly-24
AB_324004
Spleen cells from immunized mice were fused with cells of the mouse P3-NS1-1-Ag4 myeloma cell line.
Mouse anti Mouse CD44 antibody, clone 5035-41.1D recognizes human CD44, also known as Lymphocyte antigen 24, Extracellular matrix receptor III, GP90 lymphocyte homing/adhesion receptor, HUTCH-I, Phagocytic glycoprotein I (PGP-1), Hermes antigen or Hyaluronate receptor. CD44 is found on >90% bone marrow cells and spleen cells, 56% of thymus cells and 90% of lymph node cells.
Mouse anti Mouse CD44 antibody, clone 5035-41.1D recognizes the following strains of mice:- C57BL/6, C57BL/10, C57/L, C58A, AKR, 129, SJL, NZB, C3H, CE, CBA/H.
Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.
The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>).
 A, A. <i>et al.</i> (2001) Cell surface-expressed moesin-like receptor regulates T cell interactions with tissue components and binds an adhesion-modulating IL-2 peptide generated by elastase. <u>J Immunol. 166 (5): 3052-60.</u> Qian, K. <i>et al.</i> (2015) Effects of Tanshinone IIA on osteogenic differentiation of mouse bone marrow mesenchymal stem cells. <u>Naunyn Schmiedebergs Arch Pharmacol. 388 (11): 1201-9.</u>
Store at +4°C or at -20°C if preferred.
This product should be stored undiluted.
Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Health And Safety Information Regulatory		Material Safety Datasheet documentation #10041 available a https://www.bio-rad-antibodies.com/SDS/MCA1014FB 10041			t:	
		For research purposes only				
North & South America	Tel: +1 800 265 Fax: +1 919 878		Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
Email: antibody_			-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot speci	fic datashee	et for this produ	uct, please use our online : 'M364624:200529'	search tool at: bio	p-rad-antibodies.com/datashe

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint