

Datasheet: MCA1019

Description:	MOUSE ANTI HUMAN CD38
Specificity:	CD38
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AT13/5
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	TRIS buffered saline
Preservative	<0.1% sodium azide (NaN ₃)

Stabilisers

Approx. Protein Concentrations IgG concentration 1 mg/ml

Immunogen Namalwa human B-cell line.

External Database Links

UniProt:

[P28907](#) [Related reagents](#)

Entrez Gene:

[952](#) CD38 [Related reagents](#)

RRID AB_321678

Fusion Partners Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.

Specificity

Mouse anti Human CD38 antibody, clone AT13/5 recognizes CD38, also known as ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, 2'-phospho-ADP-ribosyl cyclase, 2'-phospho-ADP-ribosyl cyclase, ADP-ribosyl cyclase 1, Cyclic ADP-ribose hydrolase 1 or T10. CD38 is a 300 amino acid ~45 kDa single pass type II transmembrane glycoprotein and Mouse anti Human CD38 antibody, clone AT13/5 has a reactivity pattern consistent with other antibodies of the CD38 cluster. CD38 is expressed by plasma cells, monocytes, early lymphoid cells and activated T cells.

CD38 is widely used to study the processes of B and T differentiation and activation. CD38 is a multifunctional enzyme involved in calcium signaling and nicotinamide adenine dinucleotide (NAD⁺) metabolism ([de Oliveira et al. 2018](#)).

Flow Cytometry Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl

References

1. Corcione, A. *et al.* (2009) CX3CR1 is expressed by human B lymphocytes and mediates [corrected] CX3CL1 driven chemotaxis of tonsil centrocytes. [PLoS One. 4: e8485.](#)
2. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. [BMC Immunol. 12: 2.](#)
3. Tipping, A.J. *et al.* (2009) High GATA-2 expression inhibits human hematopoietic stem and progenitor cell function by effects on cell cycle [Blood. 113: 2661-72.](#)
4. Slyker, J.A. *et al.* (2012) Acute cytomegalovirus infection is associated with increased frequencies of activated and apoptosis-vulnerable T cells in HIV-1-infected infants. [J Virol. 86 \(20\): 11373-9.](#)
5. Slyker, J.A. *et al.* (2012) The impact of HIV-1 infection and exposure on natural killer (NK) cell phenotype in Kenyan infants during the first year of life. [Front Immunol. 3: 399.](#)
6. Kelleher, M. *et al.* (2011) Comparative Kinetics of Immune Responses and Changes in Cellular Sub-Sets Detected in Colorectal Cancer Patients Vaccinated with MVA-5T4 (TroVax) Administered Alongside Two Different Chemotherapy Regimens [J Cancer Therapy. 02 \(01\): 54-64.](#)

Further Reading	1. Malavasi, F. <i>et al.</i> (1994) Human CD38: a glycoprotein in search of a function. Immunol Today. 15 (3): 95-7.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10057 available at: https://www.bio-rad-antibodies.com/SDS/MCA101910057
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M410544:221028'

Printed on 18 Jan 2024