

## Datasheet: MCA1019

<b>Description:</b>	MOUSE ANTI HUMAN CD38
<b>Specificity:</b>	CD38
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	AT13/5
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A
<b>Buffer Solution</b>	TRIS buffered saline
<b>Preservative Stabilisers</b>	<0.1% Sodium Azide (NaN <sub>3</sub> )

<b>Approx. Protein Concentrations</b>	IgG concentration 1 mg/ml
<b>Immunogen</b>	Namalwa human B-cell line.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P28907</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">952</a>    CD38    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_321678
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD38 antibody, clone AT13/5</b> recognizes CD38, also known as ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, 2'-phospho-ADP-ribosyl cyclase, 2'-phospho-ADP-ribosyl cyclase, ADP-ribosyl cyclase 1, Cyclic ADP-ribose hydrolase 1 or T10. CD38 is a 300 amino acid ~45 kDa single pass type II transmembrane glycoprotein and Mouse anti Human CD38 antibody, clone AT13/5 has a reactivity pattern consistent with other antibodies of the CD38 cluster. CD38 is expressed by plasma cells, monocytes, early lymphoid cells and activated T cells.</p> <p>CD38 is widely used to study the processes of B and T differentiation and activation. CD38 is a multifunctional enzyme involved in calcium signaling and nicotinamide adenine dinucleotide (NAD<sup>+</sup>) metabolism (<a href="#">de Oliveira et al. 2018</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Corcione, A. <i>et al.</i> (2009) CX3CR1 is expressed by human B lymphocytes and mediates [corrected] CX3CL1 driven chemotaxis of tonsil centrocytes. <a href="#">PLoS One. 4: e8485.</a></li> <li>2. Hovden, A.O. <i>et al.</i> (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. <a href="#">BMC Immunol. 12: 2.</a></li> <li>3. Tipping, A.J. <i>et al.</i> (2009) High GATA-2 expression inhibits human hematopoietic stem and progenitor cell function by effects on cell cycle <a href="#">Blood. 113: 2661-72.</a></li> <li>4. Slyker, J.A. <i>et al.</i> (2012) Acute cytomegalovirus infection is associated with increased frequencies of activated and apoptosis-vulnerable T cells in HIV-1-infected infants. <a href="#">J Virol. 86 (20): 11373-9.</a></li> <li>5. Slyker, J.A. <i>et al.</i> (2012) The impact of HIV-1 infection and exposure on natural killer (NK) cell phenotype in Kenyan infants during the first year of life. <a href="#">Front Immunol. 3: 399.</a></li> </ol>
<b>Further Reading</b>	1. Malavasi, F. <i>et al.</i> (1994) Human CD38: a glycoprotein in search of a function. <a href="#">Immunol Today. 15 (3): 95-7.</a>
<b>Storage</b>	This product is shipped at ambient temperature. It is recommended to aliquot and store at

-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10057 available at: 10057: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10057.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10057.pdf</a>
--------------------------------------	---

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight@800</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@550</a> , <a href="#">DyLight@650</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M389268:210806'

Printed on 06 Jan 2022