

Datasheet: MCA1019PET

Description:	MOUSE ANTI HUMAN CD38:RPE
Specificity:	CD38
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	AT13/5
Isotype:	lgG1
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized			
Reconstitution	Reconstitute in 0.25 ml disilled water			
Max Ex/Em	Fluorophore Excitation Max (nm) Emission Max (nm) RPE 488nm laser 496 578			
Preparation	Purified IgG prepared by affinity chromatography on Protein A			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin			
Immunogen	Namalwa human B-cell line.			
External Database Links	UniProt: P28907 Related reagents Entrez Gene:			

CD38 Related reagents

<u>952</u>

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	Mouse anti Human CD38 antibody, clone AT13/5 recognizes a 45kDa glycoprotein and has a reactivity pattern consistent with antibodies of the CD38 cluster. CD38 is expressed by plasma cells, monocytes, early lymphoid cells and activated T cells.
	CD38 is widely used to study the processes of B and T differentiation and activation. Recently CD38 has been shown to possess ADP-ribosyl cyclase activity.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Malavasi, F. <i>et al.</i> (1994) Human CD38: a glycoprotein in search of a function. <u>Immunol Today.</u> 15 (3): 95-7.
	2. Ellis, J.H. <i>et al.</i> (1995) Engineered anti-CD38 monoclonal antibodies for immunotherapy of multiple myeloma. <u>J Immunol. 155 (2): 925-37.</u>
	3. Corcione, A. et al. (2009) CX3CR1 is expressed by human B lymphocytes and mediates
	[corrected] CX3CL1 driven chemotaxis of tonsil centrocytes. PLoS One. 4: e8485.
	4. Hovden, A.O.et al. (2011) Maturation of monocyte derived dendritic cells with OK432 boosts
	IL-12p70 secretion and conveys strong T-cell responses. BMC Immunol.;12:2.
	5. Tipping, A.J. <i>et al.</i> (2009) High GATA-2 expression inhibits human hematopoietic stem and progenitor cell function by effects on cell cycle <u>Blood. 113: 2661-72.</u>
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.
	DO NOT FREEZE.
	This product should be stored undiluted. This product is photosensitive and should be protected
	from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Email: antibody_sales_us@bio-rad.com

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America

Worldwide Fax: +1 919 878 3751

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

'M322255:180727'

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint