# Datasheet: MCA1019XZ BATCH NUMBER 163605

Description:	MOUSE ANTI HUMAN CD38:Preservative Free
Specificity:	CD38
Format:	Preservative Free
Product Type:	Monoclonal Antibody
Clone:	AT13/5
Isotype:	lgG1
Quantity:	1 mg

## **Product Details**

Applications	This product has been re	•		• • •			
	Ŭ	derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further					
	communications from the						
	information. For general	protocol r	ecommer	ndations, please visit <u>w</u>	ww.bio-		
	rad-antibodies.com/proto	rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			1/100		
	Immunohistology - Frozen	-					
	(1)	-					
	Immunohistology - Paraffin		•				
	ELISA			•			
	Immunoprecipitation	-					
	Western Blotting			•			
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as						
	•	a guide only. It is recommended that the user titrates the antibody for use in their own					
	system using appropriate negative/positive controls. (1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of						
	acetone fixation for frozen sections.						
Target Species	Human						
Product Form	Purified IgG - liquid.	Purified IgG - liquid.					
Preparation	Purified IgG prepared by affinity chromatography on Protein G.						
Buffer Solution	Phosphate buffered saline.						
Preservative	None present.						

### Stabilisers

Approx. Protein Concentrations	IgG concentration 1 mg/ml.
Immunogen	Namalwa human B-cell line.
External Database Links	UniProt: <u>P28907</u> <u>Related reagents</u> Entrez Gene:
	<u>952</u> CD38 <u>Related reagents</u>
RRID	AB_324342
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	<b>Mouse anti Human CD38 antibody, clone AT13/5</b> recognizes CD38, also known as ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, 2'-phospho-ADP-ribosyl cyclase, 2'-phospho-ADP-ribosyl cyclase, ADP-ribosyl cyclase 1, Cyclic ADP-ribose hydrolase 1 or T10. CD38 is a 300 amino acid ~45 kDa single pass type II transmembrane glycoprotein and Mouse anti Human CD38 antibody, clone AT13/5 has a reactivity pattern consistent with other antibodies of the CD38 cluster. CD38 is expressed by plasma cells, monocytes, early lymphoid cells and activated T cells.
	CD38 is widely used to study the processes of B and T differentiation and activation. CD38 is a multifunctional enzyme involved in calcium signaling and nicotinamide adenine dinucleotide (NAD+) metabolism ( <u>de Oliveira <i>et al.</i> 2018</u> ).
Flow Cytometry	Use 10µl of the suggested working dilution to label $10^6$ cells in 100µl.
References	1. Corcione, A. <i>et al.</i> (2009) CX3CR1 is expressed by human B lymphocytes and mediates [corrected] CX3CL1 driven chemotaxis of tonsil centrocytes. <u>PLoS One. 4:</u> e8485.
	<ol> <li>Hovden, A.O. <i>et al.</i> (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. <u>BMC Immunol. 12: 2.</u></li> <li>Tipping, A.J. <i>et al.</i> (2009) High GATA-2 expression inhibits human hematopoietic stem and progenitor cell function by effects on cell cycle <u>Blood. 113: 2661-72.</u></li> <li>Slyker, J.A. <i>et al.</i> (2012) Acute cytomegalovirus infection is associated with increased frequencies of activated and apoptosis-vulnerable T cells in HIV-1-infected infants. <u>J Virol. 86 (20): 11373-9.</u></li> <li>Slyker, J.A. <i>et al.</i> (2012) The impact of HIV-1 infection and exposure on natural killer (NK) cell phenotype in Kenyan infants during the first year of life. <u>Front Immunol. 3: 399.</u></li> </ol>
Further Reading	1. Malavasi, F. <i>et al.</i> (1994) Human CD38: a glycoprotein in search of a function. <u>Immunol</u> <u>Today. 15 (3): 95-7.</u>

Storage	Store at -20°C only. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA1019XZ 10162
Regulatory	For research purposes only.

## **Related Products**

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®550</u> ,			
	<u>DyLight®650</u> , <u>DyLight®680</u> , <u>DyLight®800</u> ,			
	FITC, HRP			
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>			
Goat Anti Mouse IgG (STAR77)	HRP			
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP			
Rabbit Anti Mouse IgG (STAR13)	HRP			
<b>Recommended Negative Controls</b>				

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M406894:221003'

### Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint