

Datasheet: MCA1031A647

Description:	RAT ANTI MOUSE CD45:Alexa Fluor® 647
Specificity:	CD45
Other names:	LCA
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	YW62.3
Isotype:	lgG2b
Quantity:	100 TESTS/1ml

# **Product Details**

# **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse				
Product Form	Purified IgG conjugated to Alexa Fluor 647 - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm		
	Alexa Fluor®647	650	665		
Preparation  Buffer Solution	Purified IgG prepared by affinity chromatography on Protein G from tissue supernatant  Phosphate buffered saline				
Preservative Stabilisers	0.09% sodium azide (NaN <sub>3</sub> ) 1% bovine serum albumin				
Approx. Protein Concentrations	IgG concentration 0.05	5 mg/ml			

Immunogen

Mouse spleen cells.

# **External Database**

Links

**UniProt:** 

P06800 Related reagents

**Entrez Gene:** 

19264 Ptprc Related reagents

**Synonyms** 

Ly-5

**RRID** 

AB 321156

**Fusion Partners** 

Spleen cells from immunised DA rats were fused with cells of the rat Y3/Ag1.2.3 myeloma cell line.

### **Specificity**

Rat anti Mouse CD45 antibody, clone YW62.3 recognizes the murine CD45 cell surface antigen, a single pass type1 transmembrane glycoprotein also known as protein tyrosine phosphatase receptor type C (PTPRC) and originally termed Leucocyte Common Antigen (LCA). CD45 is a 180-220kDa glycoprotein expressed by all leucocytes.

CD45 is encoded by 3 alleles in mice, differentially expressed by various inbred strains. The Ly5 gene was originally described with the gene product LY5.1 expressed in C57bl/6 and Ly5.2 expressed in SJL strains (Komura *et al.* 1975), this was subsequently expanded to include a third allele encoding Ly5.3 (Shen *et al.* 1986). Further, in 1987 a reversal of nomenclature was instigated resulting in the allele in C57bl/6 becoming Ly5<sup>b</sup> encoding Ly5.2 and the allele in SJL mice becoming Ly5<sup>a</sup> encoding Ly5.1 (Morse *et al.* 1987). Further changes were made in 1992 with Ly5.1 becoming CD45.1 (SJL) and Ly5.2 becoming CD45.2 (C57bl/6). Finally, following work demonstrating homology between the CD45 antigen and a receptor linked protein tyrosine phosphatase the CD45<sup>a</sup> gene was renamed Ptprc<sup>a</sup> and CD45<sup>b</sup> renamed Ptprc<sup>b</sup> (Charbonneau *et al.* 1988; Zebedee *et al.* 1991).

A number of different isoforms of CD45 are expressed on murine leucocytes depending on the pattern of alternative splicing of 3 exons termed A, B and C encoding regions of ~ 50 amino acids located at the N terminal region of the extracellular portion of CD45. The restricted proteins are termed CD45R with a designation depending on the expressed codon product. (Birkeland et al. 1989).

Rat anti mouse CD45 antibody, clone YW62.3 is reactive with all isoforms of murine CD45.

N.B. Some reactivity with human tissue has been observed.

# Flow Cytometry

Use 10  $\!\mu l$  of the suggested working dilution to label  $10^6$  cells in  $100 \mu l$  .

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>).

#### References

- 1. Watt, S.M. *et al.* (1987) Cell-surface markers on haemopoietic precursors. Reagents for the isolation and analysis of progenitor cell subpopulations. <u>Mol Cell Probes. 1 (4):</u> 297-326.
- 2. Kondo, Y. *et al.* (2007) Osteopetrotic (op/op) mice have reduced microglia, no Abeta deposition, and no changes in dopaminergic neurons. J Neuroinflammation. 4: 31.
- 3. Wang, S. *et al.* (2008) Drak2 contributes to West Nile virus entry into the brain and lethal encephalitis. <u>J Immunol. 181: 2084-91.</u>
- 4. Chan, D.A. *et al* (2009) Tumor vasculature is regulated by PHD2-mediated angiogenesis and bone marrow-derived cell recruitment. <u>Cancer Cell. 15: 527-38.</u>
- 5. Lee, S. *et al.* (2010) CX3CR1 deficiency alters microglial activation and reduces beta-amyloid deposition in two Alzheimer's disease mouse models. <u>Am J Pathol. 177:</u> 2549-62.
- 6. Dénes, A. *et al.* (2010) Chronic systemic infection exacerbates ischemic brain damage via a CCL5 (regulated on activation, normal T-cell expressed and secreted)-mediated proinflammatory response in mice. <u>J Neurosci. 30: 10086-95.</u>
- 7. Yoshizaki, A. *et al.* (2010) Cell adhesion molecules regulate fibrotic process via Th1/Th2/Th17 cell balance in a bleomycin-induced scleroderma model. <u>J Immunol. 185:</u> 2502-15.
- 8. Yang, J. *et al.* (2010) Evaluation of bone marrow- and brain-derived neural stem cells in therapy of central nervous system autoimmunity. <u>Am J Pathol. 177: 1989-2001.</u>
- 9. Yang, R. *et al.* (2010) Successful treatment of experimental glomerulonephritis with IdeS and EndoS, IgG-degrading streptococcal enzymes. <u>Nephrol Dial Transplant. 25:</u> 2479-86.
- 10. Reed-Geaghan, E.G. *et al.* (2010) Deletion of CD14 attenuates Alzheimer's disease pathology by influencing the brain's inflammatory milieu. J Neurosci. 30: 15369-73.
- 11. Paz, H. *et al.* (2010) The homeobox gene Hhex regulates the earliest stages of definitive hematopoiesis. <u>Blood. 116: 1254-62.</u>
- 12. Lee, D.C. *et al.* (2010) LPS- induced inflammation exacerbates phospho-tau pathology in rTg4510 mice. <u>J Neuroinflammation</u>. 7: 56.
- 13. Lebson, L. *et al.* (2010) Trafficking CD11b-positive blood cells deliver therapeutic genes to the brain of amyloid-depositing transgenic mice. <u>J Neurosci. 30: 9651-8.</u>
- 14. Long, G.G. *et al.* (2010) Hematopoietic Proliferative Lesions in the Spleen of rasH2 Transgenic Mice Treated with MNU. <u>Toxicol Pathol. 38: 1026-36.</u>
- 15. Drake, C. *et al.* (2011) Brain inflammation is induced by co-morbidities and risk factors for stroke. <u>Brain Behav Immun. 25: 1113-22.</u>
- 16. Jawhara, S. *et al.* (2012) Integrin  $\alpha X \beta_Z$  is a leukocyte receptor for *Candida albicans* and is essential for protection against fungal infections. J Immunol. 189 (5): 2468-77.
- 17. Mills, J.H. *et al.* (2012) A2A adenosine receptor signaling in lymphocytes and the central nervous system regulates inflammation during experimental autoimmune encephalomyelitis. <u>J Immunol. 188 (11): 5713-22.</u>
- 18. Zirger, J.M. *et al.* (2012) Immune-mediated loss of transgene expression from virally transduced brain cells is irreversible, mediated by IFN $\gamma$ , perforin, and TNF $\alpha$ , and due to the elimination of transduced cells. Mol Ther. 20 (4): 808-19.
- 19. Yamauchi, S. *et al.* (2012) Myosin II-dependent exclusion of CD45 from the site of Fcγ receptor activation during phagocytosis. FEBS Lett. 586: 3229-35.
- 20. Abramowski, D. *et al.* (2012) Transgenic Expression of Intraneuronal A $\beta$ 42 But Not A $\beta$ 40 Leads to Cellular A $\beta$  Lesions, Degeneration, and Functional Impairment without

Typical Alzheimer's Disease Pathology. J Neurosci. 32: 1273-83.

- 21. Chu, C.J. *et al.* (2013) Assessment and *in vivo* scoring of murine experimental autoimmune uveoretinitis using optical coherence tomography. PLoS One. 8 (5): e63002.
- 22. Murinello, S. *et al.* (2014) Fcγ receptor upregulation is associated with immune complex inflammation in the mouse retina and early age-related macular degeneration. Invest Ophthalmol Vis Sci. 55 (1): 247-58.
- 23. Yazid, S. *et al.* (2015) Annexin-A1 restricts Th17 cells and attenuates the severity of autoimmune disease. <u>J Autoimmun. 58: 1-11.</u>
- 24. Benson, C. *et al.* (2015) Voluntary wheel running delays disease onset and reduces pain hypersensitivity in early experimental autoimmune encephalomyelitis (EAE). <u>Exp.</u> Neurol. 271: 279-90.
- 25. Carbajal, K.S. *et al.* (2015) Th Cell Diversity in Experimental Autoimmune Encephalomyelitis and Multiple Sclerosis. <u>J Immunol. 195 (6): 2552-9.</u>
- 26. Kan, M.J. *et al.* (2015) Arginine deprivation and immune suppression in a mouse model of Alzheimer's disease. <u>J Neurosci. 35 (15): 5969-82.</u>
- 27. Marcos, E. *et al.* (2016) Dengue encephalitis-associated immunopathology in the mouse model: Implications for vaccine developers and antigens inducer of cellular immune response. Immunol Lett. 176: 51-6.
- 28. Haile, W.B. *et al.* (2016) The Janus kinase inhibitor ruxolitinib reduces HIV replication in human macrophages and ameliorates HIV encephalitis in a murine model. <u>Neurobiol Dis. 92 (Pt B): 137-43.</u>
- 29. Park, S.A. *et al.* (2016) Deficiency in either COX-1 or COX-2 genes does not affect amyloid beta protein burden in amyloid precursor protein transgenic mice. <u>Biochem Biophys Res Commun. 478 (1): 286-92.</u>
- 30. Srivastava, A.K. *et al.* (2016) Co-transplantation of syngeneic mesenchymal stem cells improves survival of allogeneic glial-restricted precursors in mouse brain. <u>Exp Neurol. 275</u> Pt 1: 154-61.
- 31. Zhu, C. *et al.* (2020) Antinociceptive effect of intrathecal injection of miR-9-5p modified mouse bone marrow mesenchymal stem cells on a mouse model of bone cancer pain. <u>J Neuroinflammation</u>. 17 (1): 85.
- 32. Hargreaves, A. *et al.* (2021) Tumors modulate fenestrated vascular beds and host endocrine status. J Appl Toxicol. 41 (12): 1952-65.
- 33. Filograna, R. *et al.* (2021) Mitochondrial dysfunction in adult midbrain dopamine neurons triggers an early immune response. <u>PLoS Genet. 17 (9): e1009822.</u>
- 34. Hargreaves, A. *et al.* (2022) Tumours modulate the systemic vascular response to anti-angiogenic therapy. <u>J Appl Toxicol. 42 (8): 1371-84.</u>
- 35. Chen, Y.H. *et al.* (2020) Functionally distinct IFN-γ(+) IL-17A(+) Th cells in experimental autoimmune uveitis: T-cell heterogeneity, migration, and steroid response. <u>Eur J Immunol. 50 (12): 1941-51.</u>
- 36. Lepland, A. *et al.* (2024) Therapeutic Tumor Macrophage Reprogramming in Breast Cancer Through a Peptide-Drug Conjugate <u>bioRxiv: 12 Aug [Epub ahead of print].</u>

**Storage** 

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in

frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

# Guarantee 12 months from date of despatch

# **Acknowledgements**

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

# Health And Safety Information

America

Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1031A647">https://www.bio-rad-antibodies.com/SDS/MCA1031A647</a> 10041

**Regulatory** For research purposes only

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody sales de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M407849:221010'

Fax: +44 (0)1865 852 739

## Printed on 28 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint

Fax: +1 919 878 3751