Datasheet: MCA1031G BATCH NUMBER 149480

Description:	RAT ANTI MOUSE CD45			
Specificity:	CD45			
Other names:	LCA			
Format:	Purified			
Product Type:	Monoclonal Antibody			
Clone:	YW62.3			
lsotype:	lgG2b			
Quantity:	0.25 mg			

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry				1/100 - 1/200	
	Immunohistology - Frozen					
	Immunohistology - Paraffin			•		
	ELISA			•		
	Immunoprecipitation					
	Western Blotting			•		
	Immunofluorescence	-				
	Where this antibody has not been tested for use in a particular technique this does not					
	necessarily exclude its use in such procedures. Suggested working dilutions are given as					
	a guide only. It is recommended that the user titrates the antibody for use in their own					
	0 1			•		
	system using appropriate negative/positive controls.					
Target Species	Mouse					
Product Form	Purified IgG - liquid					

Preparation	reparation Purified IgG prepared by affinity chromatography on Prote supernatant	
Buffer Solution	Phosphate buffered saline	
Preservative	0.09% Sodium Azide	

Stabilisers

Carrier Free	Yes				
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml				
Immunogen	Mouse spleen cells.				
External Database Links	UniProt: P06800 Related reagents Entrez Gene: 19264 Ptprc Related reagents				
Synonyms	Ly-5				
RRID	AB_321730				
Fusion Partners	Spleen cells from immunised DA rats were fused with cells of the rat Y3/Ag1.2.3 myeloma cell line.				
Specificity	pecificityRat anti Mouse CD45 antibody, clone YW62.3 recognizes the murine CD45 cell surface antigen, a single pass type1 transmembrane glycoprotein also known as protein tyrosine phosphatase receptor type C (PTPRC) and originally termed Leucocyte Common Antigen (LCA). CD45 is a 180-220kDa glycoprotein expressed by all leucocytes.CD45 is encoded by 3 alleles in mice, differentially expressed by various inbred strains. The Ly5 gene was originally described with the gene product LY5.1 expressed in C57bl/6 and Ly5.2 expressed in SJL strains (Komura <i>et al.</i> 1975), this was subsequently expanden to include a third allele encoding Ly5.3 (Shen <i>et al.</i> 1986). Further, in 1987 a reversal of 				
	A number of different isoforms of CD45 are expressed on murine leucocytes depending on the pattern of alternative splicing of 3 exons termed A, B and C encoding regions of ~ 50 amino acids located at the N terminal region of the extracellular portion of CD45. The restricted proteins are termed CD45R with a designation depending on the expressed codon product. (<u>Birkeland <i>et al.</i> 1989</u>).				

Rat anti mouse CD45 antibody, clone YW62.3 is reactive with all isoforms of murine CD45.

	N.B. Some reactivity with human tissue has been observed.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Watt, S.M. <i>et al.</i> (1987) Cell-surface markers on haemopoietic precursors. Reagents for the isolation and analysis of progenitor cell subpopulations. <u>Mol Cell Probes. 1 (4):</u> <u>297-326.</u> Zirger, J.M. <i>et al.</i> (2012) Immune-mediated loss of transgene expression from virally transduced brain cells is irreversible, mediated by IFNγ, perforin, and TNFα, and due to the elimination of transduced cells. <u>Mol Ther. 20 (4): 808-19.</u> Long, G.G. <i>et al.</i> (2010) Hematopoietic Proliferative Lesions in the Spleen of rasH2 Transgenic Mice Treated with MNU. <u>Toxicol Pathol. 38: 1026-36.</u>
	 Transgenic Mice Treated with MNU. <u>Toxicol Pathol. 38</u>: 1026-36. 4. Drake, C. <i>et al.</i> (2011) Brain inflammation is induced by co-morbidities and risk factors for stroke. <u>Brain Behav Immun. 25</u>: 1113-22. 5. Chan, D.A. <i>et al</i> (2009) Tumor vasculature is regulated by PHD2-mediated angiogenesis and bone marrow-derived cell recruitment. <u>Cancer Cell. 15</u>: 527-38. 6. Lebson, L. <i>et al.</i> (2010) Trafficking CD11b-positive blood cells deliver therapeutic genes to the brain of amyloid-depositing transgenic mice. <u>J Neurosci. 30</u>: 9651-8. 7. Lee, D.C. <i>et al.</i> (2010) LPS- induced inflammation exacerbates phospho-tau pathology in rTg4510 mice. <u>J Neuroinflammation. 7</u>: 56. 8. Wang, S. <i>et al.</i> (2008) Drak2 contributes to West Nile virus entry into the brain and lethal encephalitis. <u>J Immunol. 181</u>: 2084-91. 9. Paz, H. <i>et al.</i> (2010) The homeobox gene Hhex regulates the earliest stages of definitive hematopoiesis. <u>Blood. 116</u>: 1254-62. 10. Reed-Geaghan, E.G. <i>et al.</i> (2010) Deletion of CD14 attenuates Alzheimer's disease pathology by influencing the brain's inflammatory milieu. <u>J Neurosci. 30</u>: 15369-73. 11. Yang, R. <i>et al.</i> (2010) Successful treatment of experimental glomerulonephritis with IdeS and EndoS, IgG-degrading streptococcal enzymes. <u>Nephrol Dial Transplant. 25</u>: 2479-86. 12. Yang, J. <i>et al.</i> (2010) Cell adhesion molecules regulate fibrotic process via Th1/Th2/Th17 cell balance in a bleomycin-induced scleroderma model. J Immunol. 185: 2502-15. 14. Abramowski, D. <i>et al.</i> (2012) Transgenic Expression of Intraneuronal Aβ42 But Not Aβ40 Leads to Cellular Aβ Lesions, Degeneration, and Functional Impairment without Typical Alzheimer's Disease Pathology. <u>J Neurosci. 32</u>: 1273-83. 15. Dénes, A. <i>et al.</i> (2010) Chronic systemic infection exacerbates ischemic brain damage via a CCL5 (regulated on activation, normal T-cell expressed and secreted)-mediated
	 proinflammatory response in mice. J Neurosci. 30: 10086-95. 16. Kondo, Y. <i>et al.</i> (2007) Osteopetrotic (op/op) mice have reduced microglia, no Abeta deposition, and no changes in dopaminergic neurons. J Neuroinflammation. 4: 31. 17. Lee, S. <i>et al.</i> (2010) CX3CR1 deficiency alters microglial activation and reduces beta-amyloid deposition in two Alzheimer's disease mouse models. Am J Pathol. 177: 2549-62. 18. Jawhara, S. <i>et al.</i> (2012) Integrin αXβ_z is a leukocyte receptor for <i>Candida albicans</i>

and is essential for protection against fungal infections. J Immunol. 189 (5): 2468-77.

19. Yamauchi, S. *et al.* (2012) Myosin II-dependent exclusion of CD45 from the site of Fcγ receptor activation during phagocytosis. <u>FEBS Lett. 586: 3229-35.</u>

20. Yazid, S. *et al.* (2015) Annexin-A1 restricts Th17 cells and attenuates the severity of autoimmune disease. J Autoimmun. 58: 1-11.

21. Kan, M.J. *et al.* (2015) Arginine deprivation and immune suppression in a mouse model of Alzheimer's disease. <u>J Neurosci. 35 (15): 5969-82.</u>

22. Bachstetter, A.D. *et al.* (2011) Fractalkine and CX 3 CR1 regulate hippocampal neurogenesis in adult and aged rats. <u>Neurobiol Aging. 32 (11): 2030-44</u>.

23. Kuffová, L. *et al.* (2008) Cross presentation of antigen on MHC class II via the draining lymph node after corneal transplantation in mice. <u>J Immunol. 180 (3): 1353-61.</u>

24. Murinello, S. *et al.* (2014) Fcγ receptor upregulation is associated with immune complex inflammation in the mouse retina and early age-related macular degeneration. <u>Invest Ophthalmol Vis Sci. 55 (1): 247-58.</u>

25. Mills, J.H. *et al.* (2012) A2A adenosine receptor signaling in lymphocytes and the central nervous system regulates inflammation during experimental autoimmune encephalomyelitis. <u>J Immunol. 188 (11): 5713-22.</u>

26. McMorran, B.J. *et al.* (2001) G551D CF mice display an abnormal host response and have impaired clearance of *Pseudomonas* lung disease. <u>Am J Physiol Lung Cell Mol</u> <u>Physiol. 281 (3): L740-7.</u>

27. Benson, C. *et al.* (2015) Voluntary wheel running delays disease onset and reduces pain hypersensitivity in early experimental autoimmune encephalomyelitis (EAE). <u>Exp</u> Neurol. 271: 279-90.

28. Marcos, E. *et al.* (2016) Dengue encephalitis-associated immunopathology in the mouse model: Implications for vaccine developers and antigens inducer of cellular immune response. <u>Immunol Lett. 176: 51-6.</u>

29. Carbajal, K.S. *et al.* (2015) Th Cell Diversity in Experimental Autoimmune Encephalomyelitis and Multiple Sclerosis. <u>J Immunol. 195 (6): 2552-9.</u>

30. Srivastava, A.K. *et al.* (2016) Co-transplantation of syngeneic mesenchymal stem cells improves survival of allogeneic glial-restricted precursors in mouse brain. <u>Exp Neurol. 275</u> Pt 1: 154-61.

31. Haile, W.B. *et al.* (2016) The Janus kinase inhibitor ruxolitinib reduces HIV replication in human macrophages and ameliorates HIV encephalitis in a murine model. <u>Neurobiol</u> <u>Dis. 92 (Pt B): 137-43.</u>

32. Park, S.A. *et al.* (2016) Deficiency in either COX-1 or COX-2 genes does not affect amyloid beta protein burden in amyloid precursor protein transgenic mice. <u>Biochem</u> <u>Biophys Res Commun. 478 (1): 286-92.</u>

33. Chu, C.J. *et al.* (2013) Assessment and *in vivo* scoring of murine experimental autoimmune uveoretinitis using optical coherence tomography. <u>PLoS One. 8 (5): e63002.</u>

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarante	e	12 months					
Health A Informati	nd Safety on	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1031G 10040					
Regulato	ry	For research purposes only					
Relate	d Product	ts					
Recomm	Recommended Secondary Antibodies						
Rabbit A	nti Rat IgG (S	STAR16)		DyLight®800			
Rabbit A	Rabbit Anti Rat IgG (STAR17) <u>FITC</u>						
Goat Ant	Goat Anti Rat IgG (STAR72) HRP						
Goat Ant	Goat Anti Rat IgG (STAR69) FITC						
Goat Ant	Goat Anti Rat IgG (STAR73) RPE						
Rabbit A	nti Rat IgG (S	STAR21)		HRP			
Goat Ant	Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71) <u>DyLight®550</u> , <u>DyLight®650</u> , <u>DyLight®800</u>						
Goat Ant	i Rat IgG (ST	AR131)		<u>Alk. Phos., Bic</u>	otin		
North & South America	Tel: +1 800 265 7 Fax: +1 919 878 Email: antibody_s	3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio	Europe -rad.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com	
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M364663:200529'							

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint