

Datasheet: MCA1031G

BATCH NUMBER 170880

Description:	RAT ANTI MOUSE CD45
Specificity:	CD45
Other names:	LCA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	YW62.3
Isotype:	IgG2b
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% sodium azide (NaN ₃)

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Mouse spleen cells.

External Database Links

UniProt:

[P06800](#) [Related reagents](#)

Entrez Gene:

[19264](#) Ptprc [Related reagents](#)

Synonyms Ly-5

RRID AB_321730

Fusion Partners Spleen cells from immunised DA rats were fused with cells of the rat Y3/Ag1.2.3 myeloma cell line.

Specificity

Rat anti Mouse CD45 antibody, clone YW62.3 recognizes the murine CD45 cell surface antigen, a single pass type1 transmembrane glycoprotein also known as protein tyrosine phosphatase receptor type C (PTPRC) and originally termed Leucocyte Common Antigen (LCA). CD45 is a 180-220kDa glycoprotein expressed by all leucocytes.

CD45 is encoded by 3 alleles in mice, differentially expressed by various inbred strains. The Ly5 gene was originally described with the gene product LY5.1 expressed in C57bl/6 and Ly5.2 expressed in SJL strains ([Komura et al. 1975](#)), this was subsequently expanded to include a third allele encoding Ly5.3 ([Shen et al. 1986](#)). Further, in 1987 a reversal of nomenclature was instigated resulting in the allele in C57bl/6 becoming Ly5^b encoding Ly5.2 and the allele in SJL mice becoming Ly5^a encoding Ly5.1 ([Morse et al. 1987](#)). Further changes were made in 1992 with Ly5.1 becoming CD45.1 (SJL) and Ly5.2 becoming CD45.2 (C57bl/6). Finally, following work demonstrating homology between the CD45 antigen and a receptor linked protein tyrosine phosphatase the CD45^a gene was renamed Ptprc^a and CD45^b renamed Ptprc^b ([Charbonneau et al. 1988](#); [Zebedee et al. 1991](#)).

A number of different isoforms of CD45 are expressed on murine leucocytes depending on the pattern of alternative splicing of 3 exons termed A, B and C encoding regions of ~ 50 amino acids located at the N terminal region of the extracellular portion of CD45. The restricted proteins are termed CD45R with a designation depending on the expressed codon product. ([Birkeland et al. 1989](#)).

Rat anti mouse CD45 antibody, clone YW62.3 is reactive with all isoforms of murine CD45.

N.B. Some reactivity with human tissue has been observed.

Flow Cytometry Use 10 μ l of the suggested working dilution to label 10⁶ cells in 100 μ l

References

1. Watt, S.M. *et al.* (1987) Cell-surface markers on haemopoietic precursors. Reagents for the isolation and analysis of progenitor cell subpopulations. [Mol Cell Probes. 1 \(4\): 297-326.](#)
2. Kondo, Y. *et al.* (2007) Osteopetrotic (op/op) mice have reduced microglia, no A β deposition, and no changes in dopaminergic neurons. [J Neuroinflammation. 4: 31.](#)
3. Wang, S. *et al.* (2008) Drak2 contributes to West Nile virus entry into the brain and lethal encephalitis. [J Immunol. 181: 2084-91.](#)
4. Chan, D.A. *et al.* (2009) Tumor vasculature is regulated by PHD2-mediated angiogenesis and bone marrow-derived cell recruitment. [Cancer Cell. 15: 527-38.](#)
5. Lee, S. *et al.* (2010) CX3CR1 deficiency alters microglial activation and reduces beta-amyloid deposition in two Alzheimer's disease mouse models. [Am J Pathol. 177: 2549-62.](#)
6. Dénes, A. *et al.* (2010) Chronic systemic infection exacerbates ischemic brain damage via a CCL5 (regulated on activation, normal T-cell expressed and secreted)-mediated proinflammatory response in mice. [J Neurosci. 30: 10086-95.](#)
7. Yoshizaki, A. *et al.* (2010) Cell adhesion molecules regulate fibrotic process via Th1/Th2/Th17 cell balance in a bleomycin-induced scleroderma model. [J Immunol. 185: 2502-15.](#)
8. Yang, J. *et al.* (2010) Evaluation of bone marrow- and brain-derived neural stem cells in therapy of central nervous system autoimmunity. [Am J Pathol. 177: 1989-2001.](#)
9. Yang, R. *et al.* (2010) Successful treatment of experimental glomerulonephritis with IdeS and EndoS, IgG-degrading streptococcal enzymes. [Nephrol Dial Transplant. 25: 2479-86.](#)
10. Reed-Geaghan, E.G. *et al.* (2010) Deletion of CD14 attenuates Alzheimer's disease pathology by influencing the brain's inflammatory milieu. [J Neurosci. 30: 15369-73.](#)
11. Paz, H. *et al.* (2010) The homeobox gene Hhex regulates the earliest stages of definitive hematopoiesis. [Blood. 116: 1254-62.](#)
12. Lee, D.C. *et al.* (2010) LPS- induced inflammation exacerbates phospho-tau pathology in rTg4510 mice. [J Neuroinflammation. 7: 56.](#)
13. Lebson, L. *et al.* (2010) Trafficking CD11b-positive blood cells deliver therapeutic genes to the brain of amyloid-depositing transgenic mice. [J Neurosci. 30: 9651-8.](#)
14. Long, G.G. *et al.* (2010) Hematopoietic Proliferative Lesions in the Spleen of rasH2 Transgenic Mice Treated with MNU. [Toxicol Pathol. 38: 1026-36.](#)
15. Drake, C. *et al.* (2011) Brain inflammation is induced by co-morbidities and risk factors for stroke. [Brain Behav Immun. 25: 1113-22.](#)
16. Jawhara, S. *et al.* (2012) Integrin $\alpha X\beta_z$ is a leukocyte receptor for *Candida albicans* and is essential for protection against fungal infections. [J Immunol. 189 \(5\): 2468-77.](#)
17. Mills, J.H. *et al.* (2012) A2A adenosine receptor signaling in lymphocytes and the central nervous system regulates inflammation during experimental autoimmune encephalomyelitis. [J Immunol. 188 \(11\): 5713-22.](#)
18. Zirger, J.M. *et al.* (2012) Immune-mediated loss of transgene expression from virally transduced brain cells is irreversible, mediated by IFN γ , perforin, and TNF α , and due to

- the elimination of transduced cells. [Mol Ther. 20 \(4\): 808-19.](#)
19. Yamauchi, S. *et al.* (2012) Myosin II-dependent exclusion of CD45 from the site of Fcγ receptor activation during phagocytosis. [FEBS Lett. 586: 3229-35.](#)
20. Abramowski, D. *et al.* (2012) Transgenic Expression of Intraneuronal Aβ42 But Not Aβ40 Leads to Cellular Aβ Lesions, Degeneration, and Functional Impairment without Typical Alzheimer's Disease Pathology. [J Neurosci. 32: 1273-83.](#)
21. Chu, C.J. *et al.* (2013) Assessment and *in vivo* scoring of murine experimental autoimmune uveoretinitis using optical coherence tomography. [PLoS One. 8 \(5\): e63002.](#)
22. Murinello, S. *et al.* (2014) Fcγ receptor upregulation is associated with immune complex inflammation in the mouse retina and early age-related macular degeneration. [Invest Ophthalmol Vis Sci. 55 \(1\): 247-58.](#)
23. Yazid, S. *et al.* (2015) Annexin-A1 restricts Th17 cells and attenuates the severity of autoimmune disease. [J Autoimmun. 58: 1-11.](#)
24. Benson, C. *et al.* (2015) Voluntary wheel running delays disease onset and reduces pain hypersensitivity in early experimental autoimmune encephalomyelitis (EAE). [Exp Neurol. 271: 279-90.](#)
25. Carbajal, K.S. *et al.* (2015) Th Cell Diversity in Experimental Autoimmune Encephalomyelitis and Multiple Sclerosis. [J Immunol. 195 \(6\): 2552-9.](#)
26. Kan, M.J. *et al.* (2015) Arginine deprivation and immune suppression in a mouse model of Alzheimer's disease. [J Neurosci. 35 \(15\): 5969-82.](#)
27. Marcos, E. *et al.* (2016) Dengue encephalitis-associated immunopathology in the mouse model: Implications for vaccine developers and antigens inducer of cellular immune response. [Immunol Lett. 176: 51-6.](#)
28. Haile, W.B. *et al.* (2016) The Janus kinase inhibitor ruxolitinib reduces HIV replication in human macrophages and ameliorates HIV encephalitis in a murine model. [Neurobiol Dis. 92 \(Pt B\): 137-43.](#)
29. Park, S.A. *et al.* (2016) Deficiency in either COX-1 or COX-2 genes does not affect amyloid beta protein burden in amyloid precursor protein transgenic mice. [Biochem Biophys Res Commun. 478 \(1\): 286-92.](#)
30. Srivastava, A.K. *et al.* (2016) Co-transplantation of syngeneic mesenchymal stem cells improves survival of allogeneic glial-restricted precursors in mouse brain. [Exp Neurol. 275 Pt 1: 154-61.](#)
31. Zhu, C. *et al.* (2020) Antinociceptive effect of intrathecal injection of miR-9-5p modified mouse bone marrow mesenchymal stem cells on a mouse model of bone cancer pain. [J Neuroinflammation. 17 \(1\): 85.](#)
32. Hargreaves, A. *et al.* (2021) Tumors modulate fenestrated vascular beds and host endocrine status. [J Appl Toxicol. 41 \(12\): 1952-65.](#)
33. Filograna, R. *et al.* (2021) Mitochondrial dysfunction in adult midbrain dopamine neurons triggers an early immune response. [PLoS Genet. 17 \(9\): e1009822.](#)
34. Hargreaves, A. *et al.* (2022) Tumours modulate the systemic vascular response to anti-angiogenic therapy. [J Appl Toxicol. 42 \(8\): 1371-84.](#)
35. Chen, Y.H. *et al.* (2020) Functionally distinct IFN-γ(+) IL-17A(+) Th cells in experimental autoimmune uveitis: T-cell heterogeneity, migration, and steroid response. [Eur J Immunol. 50 \(12\): 1941-51.](#)
36. Lepland, A. *et al.* (2024) Therapeutic Tumor Macrophage Reprogramming in Breast Cancer Through a Peptide-Drug Conjugate [bioRxiv: 12 Aug \[Epub ahead of print\].](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1031G
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Rabbit Anti Rat IgG (STAR21...)	HRP

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M407841:221009'

Printed on 11 Mar 2026