

# Datasheet: MCA1031SBB810

Description:	RAT ANTI MOUSE CD45:StarBright Blue 810				
Specificity:	CD45				
Other names:	LCA				
Format:	StarBright Blue 810				
Product Type:	Monoclonal Antibody				
Clone:	YW62.3				
Isotype:	lgG2b				
Quantity:	100 TESTS/0.5ml				

### **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	•			Neat	
	Where this product has necessarily exclude its a guide only. It is recon system using appropria	use in such p nmended that	rocedui the use	es. Suggested workin r titrates the product f	g dilutions are given as	
Target Species	Mouse					
Product Form	Purified IgG conjugated to StarBright Blue 810 - liquid					
Max Ex/Em	Fluorophore	Excitation Max	x (nm)	Emission Max (nm)		
	StarBright Blue 810	477		802		
Preparation	Purified IgG prepared b supernatant	by affinity chro	matogr	aphy on Protein G fror	n tissue culture	
Buffer Solution	Phosphate buffered saline					
Preservative	0.09% sodium azide (NaN <sub>3</sub> )					
Stabilisers	1% bovine serum albumin					
	0.1% Pluronic F68					
	0.1% PEG 3350					
	0.05% Tween 20					

Immunogen	Mouse spleen cells.
External Database Links	UniProt: <u>P06800</u> <u>Related reagents</u> Entrez Gene: <u>19264</u> Ptprc <u>Related reagents</u>
Synonyms	Ly-5
Fusion Partners	Spleen cells from immunised DA rats were fused with cells of the rat Y3/Ag1.2.3 myeloma cell line.
Specificity	<ul> <li>Rat anti Mouse CD45 antibody, clone YW62.3 recognizes the murine CD45 cell surface antigen, a single pass type1 transmembrane glycoprotein also known as protein tyrosine phosphatase receptor type C (PTPRC) and originally termed Leucocyte Common Antigen (LCA). CD45 is a 180-220kDa glycoprotein expressed by all leucocytes.</li> <li>CD45 is encoded by 3 alleles in mice, differentially expressed by various inbred strains. The Ly5 gene was originally described with the gene product LY5.1 expressed in C57bl/6 and Ly5.2 expressed in SJL strains (Komura <i>et al.</i> 1975), this was subsequently expanded to include a third allele encoding Ly5.3 (Shen <i>et al.</i> 1986). Further, in 1987 a reversal of nomenclature was instigated resulting in the allele in C57bl/6 becoming Ly5<sup>b</sup> encoding Ly5.2 and the allele in SJL mice becoming Ly5<sup>a</sup> encoding Ly5.1 (Morse <i>et al.</i> 1987). Further changes were made in 1992 with Ly5.1 becoming CD45.1 (SJL) and Ly5.2 becoming CD45.2 (C57bl/6). Finally, following work demonstrating homology between the CD45 antigen and a receptor linked protein tyrosine phosphatase the CD45<sup>a</sup> gene was renamed Ptprc<sup>a</sup> and CD45<sup>b</sup> renamed Ptprc<sup>b</sup> (Charbonneau <i>et al.</i> 1988; Zebedee <i>et al.</i> 1991).</li> <li>A number of different isoforms of CD45 are expressed on murine leucocytes depending an the pattern of alternative enlicing of 2 event termed A. B. and C. expending regione of were series of a straine of alternative enlicing of 2 event termed A. B. and C. expending regione of were series of a straine of alternative enlicing of 2 event termed A. B. and C.</li> </ul>
	on the pattern of alternative splicing of 3 exons termed A, B and C encoding regions of ~ 50 amino acids located at the N terminal region of the extracellular portion of CD45. The restricted proteins are termed CD45R with a designation depending on the expressed codon product. ( <u>Birkeland <i>et al.</i> 1989</u> ). Rat anti mouse CD45 antibody, clone YW62.3 is reactive with all isoforms of murine
	CD45.
	N.B. Some reactivity with human tissue has been observed.
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	1. Watt, S.M. <i>et al.</i> (1987) Cell-surface markers on haemopoietic precursors. Reagents for the isolation and analysis of progenitor cell subpopulations. <u>Mol Cell Probes. 1 (4):</u> <u>297-326.</u>

Kondo, Y. *et al.* (2007) Osteopetrotic (op/op) mice have reduced microglia, no Abeta deposition, and no changes in dopaminergic neurons. <u>J Neuroinflammation. 4: 31.</u>
 Wang, S. *et al.* (2008) Drak2 contributes to West Nile virus entry into the brain and lethal encephalitis. <u>J Immunol. 181: 2084-91.</u>

4. Chan, D.A. *et al* (2009) Tumor vasculature is regulated by PHD2-mediated angiogenesis and bone marrow-derived cell recruitment. <u>Cancer Cell. 15: 527-38.</u>
5. Lee, S. *et al.* (2010) CX3CR1 deficiency alters microglial activation and reduces beta-amyloid deposition in two Alzheimer's disease mouse models. <u>Am J Pathol. 177:</u> 2549-62.

6. Dénes, A. *et al.* (2010) Chronic systemic infection exacerbates ischemic brain damage via a CCL5 (regulated on activation, normal T-cell expressed and secreted)-mediated proinflammatory response in mice. <u>J Neurosci. 30: 10086-95.</u>

7. Yoshizaki, A. *et al.* (2010) Cell adhesion molecules regulate fibrotic process via Th1/Th2/Th17 cell balance in a bleomycin-induced scleroderma model. <u>J Immunol. 185:</u> <u>2502-15.</u>

8. Yang, J. *et al.* (2010) Evaluation of bone marrow- and brain-derived neural stem cells in therapy of central nervous system autoimmunity. <u>Am J Pathol. 177: 1989-2001.</u>

9. Yang, R. *et al.* (2010) Successful treatment of experimental glomerulonephritis with IdeS and EndoS, IgG-degrading streptococcal enzymes. <u>Nephrol Dial Transplant. 25:</u> 2479-86.

Reed-Geaghan, E.G. *et al.* (2010) Deletion of CD14 attenuates Alzheimer's disease pathology by influencing the brain's inflammatory milieu. <u>J Neurosci. 30: 15369-73.</u>
 Paz, H. *et al.* (2010) The homeobox gene Hhex regulates the earliest stages of definitive hematopoiesis. <u>Blood. 116: 1254-62.</u>

12. Lee, D.C. *et al.* (2010) LPS- induced inflammation exacerbates phospho-tau pathology in rTg4510 mice. J Neuroinflammation. 7: 56.

13. Lebson, L. *et al.* (2010) Trafficking CD11b-positive blood cells deliver therapeutic genes to the brain of amyloid-depositing transgenic mice. <u>J Neurosci. 30: 9651-8.</u>

14. Long, G.G. *et al.* (2010) Hematopoietic Proliferative Lesions in the Spleen of rasH2 Transgenic Mice Treated with MNU. <u>Toxicol Pathol. 38: 1026-36.</u>

15. Drake, C. *et al.* (2011) Brain inflammation is induced by co-morbidities and risk factors for stroke. <u>Brain Behav Immun. 25: 1113-22.</u>

16. Jawhara, S. *et al.* (2012) Integrin  $\alpha X \beta_z$  is a leukocyte receptor for *Candida albicans* and is essential for protection against fungal infections. J Immunol. 189 (5): 2468-77.

17. Mills, J.H. *et al.* (2012) A2A adenosine receptor signaling in lymphocytes and the central nervous system regulates inflammation during experimental autoimmune encephalomyelitis. <u>J Immunol. 188 (11): 5713-22.</u>

18. Zirger, J.M. *et al.* (2012) Immune-mediated loss of transgene expression from virally transduced brain cells is irreversible, mediated by IFN $\gamma$ , perforin, and TNF $\alpha$ , and due to the elimination of transduced cells. <u>Mol Ther. 20 (4): 808-19.</u>

19. Yamauchi, S. *et al.* (2012) Myosin II-dependent exclusion of CD45 from the site of Fcγ receptor activation during phagocytosis. <u>FEBS Lett. 586: 3229-35.</u>

20. Abramowski, D. *et al.* (2012) Transgenic Expression of Intraneuronal A $\beta$ 42 But Not A $\beta$ 40 Leads to Cellular A $\beta$  Lesions, Degeneration, and Functional Impairment without Typical Alzheimer's Disease Pathology. <u>J Neurosci. 32: 1273-83.</u>

21. Chu, C.J. *et al.* (2013) Assessment and *in vivo* scoring of murine experimental autoimmune uveoretinitis using optical coherence tomography. <u>PLoS One. 8 (5): e63002.</u>

	22. Murinello, S. <i>et al.</i> (2014) Fcγ receptor upregulation is associated with immune complex inflammation in the mouse retina and early age-related macular degeneration. Invest Ophthalmol Vis Sci. 55 (1): 247-58.
	23. Yazid, S. <i>et al.</i> (2015) Annexin-A1 restricts Th17 cells and attenuates the severity of
	autoimmune disease. <u>J Autoimmun. 58: 1-11.</u>
	24. Benson, C. et al. (2015) Voluntary wheel running delays disease onset and reduces
	pain hypersensitivity in early experimental autoimmune encephalomyelitis (EAE). Exp
	Neurol. 271: 279-90.
	25. Carbajal, K.S. et al. (2015) Th Cell Diversity in Experimental Autoimmune
	Encephalomyelitis and Multiple Sclerosis. <u>J Immunol. 195 (6): 2552-9.</u>
	26. Kan, M.J. et al. (2015) Arginine deprivation and immune suppression in a mouse
	model of Alzheimer's disease. <u>J Neurosci. 35 (15): 5969-82.</u>
	27. Marcos, E. et al. (2016) Dengue encephalitis-associated immunopathology in the
	mouse model: Implications for vaccine developers and antigens inducer of cellular
	immune response. Immunol Lett. 176: 51-6.
	28. Haile, W.B. et al. (2016) The Janus kinase inhibitor ruxolitinib reduces HIV replication
	in human macrophages and ameliorates HIV encephalitis in a murine model. <u>Neurobiol</u>
	<u>Dis. 92 (Pt B): 137-43.</u>
	29. Park, S.A. et al. (2016) Deficiency in either COX-1 or COX-2 genes does not affect
	amyloid beta protein burden in amyloid precursor protein transgenic mice. <u>Biochem</u>
	<u>Biophys Res Commun. 478 (1): 286-92.</u>
	30. Srivastava, A.K. et al. (2016) Co-transplantation of syngeneic mesenchymal stem cells
	improves survival of allogeneic glial-restricted precursors in mouse brain. <u>Exp Neurol. 275</u> Pt 1: 154-61.
	31. Zhu, C. <i>et al.</i> (2020) Antinociceptive effect of intrathecal injection of miR-9-5p modified mouse bone marrow mesenchymal stem cells on a mouse model of bone cancer pain. $\underline{J}$ Neuroinflammation. 17 (1): 85.
	32. Hargreaves, A. <i>et al.</i> (2021) Tumors modulate fenestrated vascular beds and host
	endocrine status. J Appl Toxicol. 41 (12): 1952-65.
	33. Filograna, R. <i>et al.</i> (2021) Mitochondrial dysfunction in adult midbrain dopamine
	neurons triggers an early immune response. <u>PLoS Genet. 17 (9): e1009822.</u>
	34. Hargreaves, A. <i>et al.</i> (2022) Tumours modulate the systemic vascular response to
	anti-angiogenic therapy. <u>J Appl Toxicol. 42 (8): 1371-84.</u>
	35. Chen, Y.H. <i>et al.</i> (2020) Functionally distinct IFN- $\gamma$ (+) IL-17A(+) Th cells in
	experimental autoimmune uveitis: T-cell heterogeneity, migration, and steroid response. Eur J Immunol. 50 (12): 1941-51.
Storage	Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1031SBB810 20471

## Related Products Recommended Useful Reagents

#### MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21		
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50		
	Email: antibody_sales_us@bio	o-rad.com	Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com		
To find a batch/lat apositic detection for this product, places use our opling search tool at: his red optibadies com/detections							

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M411600:221104'

#### Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint