

Datasheet: MCA1033F

BATCH NUMBER 155159

Description:	RAT ANTI MOUSE CD71:FITC
Specificity:	CD71
Other names:	TRANSFERRIN RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	YTA74.4
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by ion exchange chromatography		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Concanavalin A activated mouse spleen cells.
External Database Links	<p>UniProt: Q62351 Related reagents</p> <p>Entrez Gene: 22042 Tfrc Related reagents</p>
Synonyms	Tfrr
RRID	AB_321759
Fusion Partners	Spleen cells from an immunised DA rat were fused with cells of the Y3/Ag1.2.3 rat myeloma cell line.
Specificity	<p>Rat anti Mouse CD71 antibody, clone YTA74.4 recognizes the mouse Transferrin receptor protein 1 also known as CD71 or TfR1. CD71 is a 763 amino acid glycoprotein homodimer of ~95 kDa subunits. CD71 is expressed by dividing cells, and functions as a transferrin receptor mediating uptake of iron.</p> <p>Rat anti Mouse CD71 antibody, clone YTA74.4 blocks the binding of R17 217.1.3. and R17 208.2 anti-TFR monoclonal antibodies (Trowbridge et al. 1982).</p>
Flow Cytometry	<p>Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
References	<ol style="list-style-type: none"> 1. Millot, S. <i>et al.</i> (2010) Erythropoietin stimulates spleen BMP4-dependent stress erythropoiesis and partially corrects anemia in a mouse model of generalized inflammation. Blood. 116: 6072-81. 2. Kuo, Y.M. <i>et al.</i> (2004) Mislocalisation of hephaestin, a multicopper ferroxidase involved in basolateral intestinal iron transport, in the sex linked anaemia mouse. Gut. 53: 201-6. 3. Krysiak, K. <i>et al.</i> (2014) Reduced Levels of Hspa9 Attenuates Stat5 Activation in Mouse B-cells. Exp Hematol. pii: S0301-472X(14)00817-0. 4. Byun, M. <i>et al.</i> (2007) Cowpox virus exploits the endoplasmic reticulum retention pathway to inhibit MHC class I transport to the cell surface. Cell Host Microbe. 2: 306-15. 5. Ripich, T. and Jessberger, R. (2011) SWAP-70 regulates erythropoiesis by controlling α4 integrin. Haematologica. 96: 1743-52. 6. Hadziahmetovic, M. <i>et al.</i> (2012) Microarray analysis of murine retinal light damage reveals changes in iron regulatory, complement, and antioxidant genes in the neurosensory retina and isolated RPE. Invest Ophthalmol Vis Sci. 53 (9): 5231-41. 7. Niewoehner, J. <i>et al.</i> (2014) Increased brain penetration and potency of a therapeutic antibody using a monovalent molecular shuttle. Neuron. 81: 49-60.

8. Sands, S.A. *et al.* (2015) The habenula and iron metabolism in cerebral mouse models of multiple sclerosis. [Neurosci Lett. 606: 204-8.](#)
9. Baumann, B. *et al.* (2017) Conditional Müller Cell Ablation Leads to Retinal Iron Accumulation. [Invest Ophthalmol Vis Sci. 58 \(10\): 4223-34.](#)
10. Nelvagal, H.R. *et al.* (2020) Comparative proteomic profiling reveals mechanisms for early spinal cord vulnerability in CLN1 disease. [Sci Rep. 10 \(1\): 15157.](#)

Further Reading

1. Lesley, J. *et al.* (1984) Expression of transferrin receptor on murine hematopoietic progenitors. [Cell Immunol. 83 \(1\): 14-25.](#)
2. Trowbridge, I.S. *et al.* (1982) Murine cell surface transferrin receptor: studies with an anti-receptor monoclonal antibody. [J Cell Physiol. 112 \(3\): 403-10.](#)

Storage

Store at +4°C for one month or at -20°C for longer.

This product should be stored undiluted.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1033F>
10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M364671:200529'

Printed on 15 Apr 2024