

Datasheet: MCA1033F

Description:	RAT ANTI MOUSE CD71:FITC
Specificity:	CD71
Other names:	TRANSFERRIN RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	YTA74.4
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Concanavalin A activated mouse spleen cells.
External Database Links	<p>UniProt: Q62351 Related reagents</p> <p>Entrez Gene: 22042 Tfrc Related reagents</p>
Synonyms	Tfrr
RRID	AB_321759
Fusion Partners	Spleen cells from an immunized DA rat were fused with cells of the Y3/Ag1.2.3 rat myeloma cell line.
Specificity	<p>Rat anti Mouse CD71 antibody, clone YTA74.4 recognizes the mouse transferrin receptor protein 1 also known as CD71 or TfR1. CD71 is a 763 amino acid glycoprotein homodimer of ~95 kDa subunits. CD71 is expressed by dividing cells, and functions as a transferrin receptor mediating uptake of iron.</p> <p>Rat anti Mouse CD71 antibody, clone YTA74.4 blocks the binding of R17 217.1.3. and R17 208.2 anti-TFR monoclonal antibodies (Trowbridge et al. 1982).</p>
Flow Cytometry	<p>Use 10up of the suggested working dilution to label 10⁶ cells in 100µl.</p> <p>The Fc region of monoclonal antibodies may bind to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
References	<ol style="list-style-type: none"> 1. Millot, S. <i>et al.</i> (2010) Erythropoietin stimulates spleen BMP4-dependent stress erythropoiesis and partially corrects anemia in a mouse model of generalized inflammation. Blood. 116: 6072-81. 2. Kuo, Y.M. <i>et al.</i> (2004) Mislocalisation of hephaestin, a multicopper ferroxidase involved in basolateral intestinal iron transport, in the sex linked anaemia mouse. Gut. 53: 201-6. 3. Krysiak, K. <i>et al.</i> (2015) Reduced levels of Hspa9 attenuate Stat5 activation in mouse B cells. Exp Hematol. 43 (4): 319-30.e10. 4. Byun, M. <i>et al.</i> (2007) Cowpox virus exploits the endoplasmic reticulum retention pathway to inhibit MHC class I transport to the cell surface. Cell Host Microbe. 2: 306-15. 5. Ripich, T. and Jessberger, R. (2011) SWAP-70 regulates erythropoiesis by controlling α4 integrin. Haematologica. 96: 1743-52. 6. Hadziahmetovic, M. <i>et al.</i> (2012) Microarray analysis of murine retinal light damage reveals changes in iron regulatory, complement, and antioxidant genes in the neurosensory retina and isolated RPE. Invest Ophthalmol Vis Sci. 53 (9): 5231-41. 7. Niewoehner, J. <i>et al.</i> (2014) Increased brain penetration and potency of a therapeutic antibody using a monovalent molecular shuttle. Neuron. 81: 49-60. 8. Sands, S.A. <i>et al.</i> (2015) The habenula and iron metabolism in cerebral mouse models

- of multiple sclerosis. [Neurosci Lett. 606: 204-8.](#)
9. Baumann, B. *et al.* (2017) Conditional Müller Cell Ablation Leads to Retinal Iron Accumulation. [Invest Ophthalmol Vis Sci. 58 \(10\): 4223-34.](#)
 10. Nelvagal, H.R. *et al.* (2020) Comparative proteomic profiling reveals mechanisms for early spinal cord vulnerability in CLN1 disease. [Sci Rep. 10 \(1\): 15157.](#)
 11. Hargreaves, A. *et al.* (2021) Tumors modulate fenestrated vascular beds and host endocrine status. [J Appl Toxicol. 41 \(12\): 1952-65.](#)
 12. Zhang, K.R. *et al.* (2022) Conditional knockout of hephaestin in the neural retina disrupts retinal iron homeostasis. [Exp Eye Res. 218: 109028.](#)
 13. Hargreaves, A. *et al.* (2022) Tumours modulate the systemic vascular response to anti-angiogenic therapy. [J Appl Toxicol. 42 \(8\): 1371-84.](#)
 14. Hargreaves, A. *et al.* (2021) Tumors modulate fenestrated vascular beds and host endocrine status. [J Appl Toxicol. 41 \(12\): 1952-65.](#)

Further Reading	<ol style="list-style-type: none"> 1. Lesley, J. <i>et al.</i> (1984) Expression of transferrin receptor on murine hematopoietic progenitors. Cell Immunol. 83 (1): 14-25. 2. Trowbridge, I.S. <i>et al.</i> (1982) Murine cell surface transferrin receptor: studies with an anti-receptor monoclonal antibody. J Cell Physiol. 112 (3): 403-10.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1033F10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M438599:250520'

Printed on 23 May 2025