

Datasheet: MCA1033GA

## **BATCH NUMBER 1804**

Description:	RAT ANTI MOUSE CD71	
Specificity:	CD71	
Other names:	TRANSFERRIN RECEPTOR	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	YTA74.4	
Isotype:	lgG2a	
Quantity:	0.1 mg	

# **Product Details**

# **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Immunohistology - Frozen	•			
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation	•			
Western Blotting			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein (supernatant	G from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide	

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Concanavilin A activated mouse spleen cells.
External Database Links	UniProt:  Q62351 Related reagents  Entrez Gene:
	22042 Tfrc Related reagents
Synonyms	Trfr
RRID	AB_324703
Fusion Partners	Spleen cells from an immunised DA rat were fused with cells of the Y3/Ag1.2.3 rat myeloma cell line.
Specificity	Rat anti Mouse CD71 antibody, clone YTA74.4 recognizes the mouse Transferrin receptor protein 1 also known as CD71 or TfR1. CD71 is a 763 amino acid glycoprotein homodimer of ~95 kDa subunits. CD71 is expressed by dividing cells, and functions as a transferrin receptor mediating uptake of iron.  Rat anti Mouse CD71 antibody, clone YTA74.4 blocks the binding of R17 217.1.3. and R17 208.2 anti-TFR monoclonal antibodies (Trowbridge et al. 1982).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1 x $10^6$ cells in 100ul.
References	<ol> <li>Millot, S. <i>et al.</i> (2010) Erythropoietin stimulates spleen BMP4-dependent stress erythropoiesis and partially corrects anemia in a mouse model of generalized inflammation. Blood. 116: 6072-81.</li> <li>Kuo, Y.M. <i>et al.</i> (2004) Mislocalisation of hephaestin, a multicopper ferroxidase involved in basolateral intestinal iron transport, in the sex linked anaemia mouse. Gut. 53: 201-6.</li> <li>Krysiak, K. <i>et al.</i> (2014) Reduced Levels of Hspa9 Attenuates Stat5 Activation in Mouse B-cells. Exp Hematol. pii: S0301-472X(14)00817-0.</li> <li>Byun, M. <i>et al.</i> (2007) Cowpox virus exploits the endoplasmic reticulum retention pathway to inhibit MHC class I transport to the cell surface. Cell Host Microbe. 2: 306-15.</li> <li>Ripich, T. and Jessberger, R. (2011) SWAP-70 regulates erythropoiesis by controlling α4 integrin. Haematologica. 96: 1743-52.</li> <li>Hadziahmetovic, M. <i>et al.</i> (2012) Microarray analysis of murine retinal light damage reveals changes in iron regulatory, complement, and antioxidant genes in the neurosensory retina and isolated RPE. Invest Ophthalmol Vis Sci. 53 (9): 5231-41.</li> <li>Niewoehner, J. <i>et al.</i> (2014) Increased brain penetration and potency of a therapeutic antibody using a monovalent molecular shuttle. Neuron. 81: 49-60.</li> <li>Sands, S.A. <i>et al.</i> (2015) The habenula and iron metabolism in cerebral mouse models</li> </ol>

of multiple sclerosis. Neurosci Lett. 606: 204-8.

9. Baumann, B. *et al.* (2017) Conditional Müller Cell Ablation Leads to Retinal Iron Accumulation. Invest Ophthalmol Vis Sci. 58 (10): 4223-34.

10. Nelvagal, H.R. *et al.* (2020) Comparative proteomic profiling reveals mechanisms for early spinal cord vulnerability in CLN1 disease. Sci Rep. 10 (1): 15157.

### **Further Reading**

- 1. Lesley, J. *et al.* (1984) Expression of transferrin receptor on murine hematopoietic progenitors. Cell Immunol. 83 (1): 14-25.
- 2. Trowbridge, I.S. *et al.* (1982) Murine cell surface transferrin receptor: studies with an anti-receptor monoclonal antibody. <u>J Cell Physiol. 112 (3): 403-10.</u>

### Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

# Guarantee

12 months from date of despatch

# Health And Safety Information

Material Safety Datasheet documentation #10040 available at:

https://www.bio-rad-antibodies.com/SDS/MCA1033GA

10040

Regulatory

For research purposes only

## Related Products

### **Recommended Secondary Antibodies**

Rabbit Anti Rat IgG (STAR16...) <u>DyLight®800</u>

Rabbit Anti Rat IgG (STAR17...)

Goat Anti Rat IgG (STAR72...)

HRP

Goat Anti Rat IgG (STAR69...)

Goat Anti Rat IgG (STAR73...)

RPE

Rabbit Anti Rat IgG (STAR21...)

HRP

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) DyLight®550, DyLight®650, DyLight®800

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

### **Recommended Negative Controls**

### RAT IgG2a NEGATIVE CONTROL (MCA1212)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M364673:200529'

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint