Datasheet: MCA1037APC BATCH NUMBER 1701

Description:	RAT ANTI DOG CD5:APC			
Specificity:	CD5			
Format:	APC			
Product Type:	Monoclonal Antibody			
Clone:	YKIX322.3			
Isotype:	lgG2a			
Quantity:	100 TESTS			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			Neat		
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.						
Target Species	Dog						
Product Form	Purified IgG conjugated to Allophycocyanin (APC) - lyophilised						
Reconstitution	Reconstitute with 1ml distilled water						
Max Ex/Em	Fluorophore	Excitation Max	c (nm) E	mission Max (nm)			
	APC	650		661			
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 5% Sucrose						

Immunogen	Concanavilin A activated canine peripheral blood cells
RRID	AB_1833728
Fusion Partners	Spleen cells from an immunised DA rat were fused with cells of the rat Y3/Ag1.2.3 myeloma cell line
Specificity	Rat anti Dog CD5 antibody, clone YKIX322.3 recognizes canine CD5, a 67 kDa cell surface type 1 transmembrane glycoprotein also known as lymphocyte antigen T1, Ly-1 or Leu-1. CD5 is expressed on the surface of T-cells and thymocytes, CD5 is also expressed by NK cells at low levels (<u>Huang <i>et al.</i> 2008</u>). Rat anti dog CD5, cloneYKIX322.3 was clustered as canine CD5 in the First Canine Leucocyte Antigen Workshop (<u>Cobbold <i>et al.</i></u> 1994).
	In a study of 73 cases of canine chronic lymphocytic leukemia (CLL) CD5 expression was absent on all cases of B-cell CLL as defined by CD21 expression and lack of CD3 or other T cell antigen expression (<u>Vernau and Moore 1999</u>). Rat anti dog CD5 serves as a useful marker for the discrimination of canine leukemias of differing origins (<u>Deravi <i>et al.</i> 2017</u>).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Cobbold, S/ & Metcalfe, S. (1994) Monoclonal antibodies that define canine homologues of human CD antigens: summary of the First International Canine Leukocyte Antigen Workshop (CLAW). <u>Tissue Antigens. 43 (3): 137-54.</u> Hewicker-Trautwein, M. <i>et al.</i> (1999) Immunocytochemical demonstration of lymphocyte subsets and MHC class II antigen expression in synovial membranes from dogs with rheumatoid arthritis and degenerative joint disease. <u>Vet Immunol Immunopathol. 67 (4):</u> <u>341-57.</u> Huang, Y.C. (2008) CD5-low expression lymphocytes in canine peripheral blood show characteristics of natural killer cells. <u>J Leukoc Biol. 84: 1501-10.</u> Araújo, M.S. <i>et al.</i> (2011) Immunological changes in canine peripheral blood leukocytes triggered by immunization with first or second generation vaccines against canine visceral leishmaniasis. <u>Vet Immunol Immunopathol. 141: 64-75.</u> Burnett, R.C. <i>et al.</i> (2003) Diagnosis of canine lymphoid neoplasia using clonal rearrangements of antigen receptor genes. <u>Vet Pathol. 40: 32-41.</u> Fosmire, S.P. <i>et al.</i> (2007) Inactivation of the p16 cyclin-dependent kinase inhibitor in high-grade canine non-Hodgkin's T-cell lymphoma. <u>Vet Pathol. 44: 467-78.</u> Guarga, J.L. <i>et al.</i> (2002) Evaluation of a specific immunochemotherapy for the treatment of canine visceral leishmaniasis. <u>Vet Immunol Immunopathol. 88: 13-20.</u> Vernau, W. Moore, P.F. <i>et al.</i> (1999) An immunophenotypic study of canine leukemias and preliminary assessment of clonality by polymerase chain reaction. <u>Vet Immunol Immunopathol. 69: 145-64.</u> Lamerato-kozicki, A.R. <i>et al.</i> (2006) Canine hemangiosarcoma originates from hematopoietic precursors with potential for endothelial differentiation. <u>Exp Hematol. 34 (7): 870-8.</u> Rütgen BC <i>et al.</i> (2012) Authentication of primordial characteristics of the CLBL-1 cell line prove the integrity of a canine B-cell lymphoma in a murine in vivo model. <u>PLoS One.</u> <u>7 (6</u>

11. Aresu, L. *et al.* (2014) VEGF and MMP-9: biomarkers for canine lymphoma. <u>Vet Comp</u> <u>Oncol. 12: 29-36.</u>

12. Moreira, M.L. *et al.* (2016) Vaccination against canine leishmaniosis increases the phagocytic activity, nitric oxide production and expression of cell activation/migration molecules in neutrophils and monocytes. <u>Vet Parasitol. 220: 33-45.</u>

13. Gelain, M.E. *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. <u>Vet Immunol Immunopathol. 159 (1-2): 91-6.</u>

14. Michael, H.T. *et al.* (2013) Isolation and characterization of canine natural killer cells. <u>Vet Immunol Immunopathol. 155 (3): 211-7.</u>

15. Karayannopoulou, M. *et al.* (2017) Evaluation of blood T-lymphocyte subpopulations involved in host cellular immunity in dogs with mammary cancer. <u>Vet Immunol Immunopathol. 186: 45-50.</u>

16. Bonnefont-Rebeix, C. *et al.* (2016) Characterization of a novel canine T-cell line established from a spontaneously occurring aggressive T-cell lymphoma with large granular cell morphology. <u>Immunobiology. 221 (1): 12-22.</u>

17. Gibbons, N. *et al.* (2017) Phenotypic heterogeneity of peripheral monocytes in healthy dogs. <u>Vet Immunol Immunopathol. 190: 26-30.</u>

18. Deravi, N. *et al.* (2017) Specific immunotypes of canine T cell lymphoma are associated with different outcomes. <u>Vet Immunol Immunopathol. 191: 5-13.</u>

19. GomesMde, O. *et al.* (2011) Old beagle dogs have lower faecal concentrations of some fermentation products and lower peripheral lymphocyte counts than young adult beagles. <u>Br J Nutr. 106 Suppl 1: S187-90.</u>

20. MariaA, P.J. *et al.* (2017) The effect of age and carbohydrate and protein sources on digestibility, fecal microbiota, fermentation products, fecal IgA, and immunological blood parameters in dogs. <u>J Anim Sci. 95 (6): 2452-66.</u>

21. Stokol, T. *et al.* (2015) Alkaline phosphatase is a useful cytochemical marker for the diagnosis of acute myelomonocytic and monocytic leukemia in the dog. <u>Vet Clin Pathol.</u> <u>44 (1): 79-93.</u>

22. Martini, V. *et al.* (2019) Prognostic role of non-neoplastic lymphocytes in lymph node aspirates from dogs with diffuse large B-cell lymphoma treated with chemo-immunotherapy. <u>Res Vet Sci. 125: 130-5.</u>

23. Lin, C.S. *et al.* (2018) Activating natural killer (NK) cytotoxicity of canine CD5⁻CD21⁻ cells requires low surface CD5 density NK cells. <u>Iran J Vet Res. 19 (2): 87-95.</u>

24. Roatt, B.M. *et al.* (2017) A Vaccine Therapy for Canine Visceral Leishmaniasis Promoted Significant Improvement of Clinical and Immune Status with Reduction in Parasite Burden. <u>Front Immunol. 8: 217.</u>

25. Aricò, A. *et al.* (2013) The role of vascular endothelial growth factor and matrix metalloproteinases in canine lymphoma: *in vivo* and *in vitro* study. <u>BMC Vet Res. 9: 94.</u>
26. Aguiar-Soares, R.D.O. *et al.* (2020) Phase I and II Clinical Trial Comparing the LBSap, Leishmune[®], and Leish-Tec[®] Vaccines against Canine Visceral Leishmaniasis. <u>Vaccines</u> (Basel). 8 (4)Nov 17 [Epub ahead of print].

27. Graves, S.S. *et al.* (2019) Development and characterization of a canine-specific anti-CD94 (KLRD-1) monoclonal antibody. <u>Vet Immunol Immunopathol. 211: 10-8.</u>
28. Ito, D. *et al.* (2015) A double blinded, placebo-controlled pilot study to examine reduction of CD34 ⁺/CD117 ⁺/CD133 ⁺ lymphoma progenitor cells and duration of remission induced by neoadjuvant valspodar in dogs with large B-cell lymphoma. <u>F1000Res. 4: 42.</u>

	 29. Wolf-Ringwall, A. <i>et al.</i> (2020) Prospective evaluation of flow cytometric characteristics, histopathologic diagnosis and clinical outcome in dogs with naïve B-cell lymphoma treated with a 19-week CHOP protocol. <u>Vet Comp Oncol. 18 (3): 342-52.</u> 30. Sayag, D. <i>et al.</i> (2020) Proof-of-concept study: Evaluation of plasma and urinary electrolytes as markers of response to L-asparaginase therapy in dogs with high-grade
	lymphoma. <u>Vet Clin Pathol. 49 (3): 476-83.</u>
Storage	Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA1037APC 20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

RAT IgG2a NEGATIVE CONTROL:APC (MCA6005APC)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50		
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com		
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375245:210104'							

Printed on 05 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint