Datasheet: MCA1039PB BATCH NUMBER 150386

Description: RAT ANTI DOG CD8:Pacific		
Specificity:	CD8	
Format:	Pacific Blue®	
Product Type:	Monoclonal Antibody	
Clone:	YCATE55.9	
lsotype:	lgG1	
Quantity:	100 TESTS/1ml	

Product Details

Applications	This product has been derived from testing w communications from information. For gener rad-antibodies.com/pro	ithin our labora the originators al protocol rec	atories, p s. Please	peer-reviewed publicate refer to references in	tions or personal dicated for further
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	-			Neat
	Where this antibody have			•	•
	necessarily exclude its a guide only. It is reco systems with appropria	mmended that	t the use	r titrates the antibody	g dilutions are given as for use in their own
Target Species	Dog				
Product Form	Purified IgG conjugate	ed to Pacific Bl	ue® - liq	luid	
Max Ex/Em	Fluorophore	Excitation Ma	ıx (nm)	Emission Max (nm)	
	Pacific Blue®	410		455	
Preparation	Purified IgG prepared supernatant	by affinity chro	omatogra	aphy on Protein G fron	n tissue culture
Buffer Solution	Phosphate buffered sa	aline			
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum /	Albumin			
Approx. Protein Concentrations	IgG concentration 0.05	5 mg/ml			

External Database Links	UniProt: P33706 Related reagents
	Entrez Gene:
	403157 CD8A Related reagents
RRID	AB_1102342
Fusion Partners	Spleen cells from immunised DA rat were fused with cells of the Y3/Ag1.2.3 rat myeloma cell line.
Specificity	Rat anti Dog CD8 antibody, clone YCATE55.9 was clustered as Canine CD8 in the First Canine Leukocyte Antigen Workshop (<u>Cobbold <i>et al.</i> 1994</u>). YCATE55.9 reacts with a rat cell line transfected with cDNA for canine CD8 α (<u>Gorman <i>et al.</i> 1994</u>) and blocks MHC class I dependent T-cell responses <i>in vitro</i> and <i>in vivo</i> .
	Rat anti Dog CD8, clone YCATE55.9 has been shown to deplete circulating CD8+ T cells when administered to dogs <i>in vivo</i> . (Watson <i>et al.</i> 1993) Reduced levels of circulating CD8+ T cells has been associated with decreased survival times for dogs with osteosarcoma (Biller <i>et al.</i> 2010).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Cobbold, S. & Metcalfe, S. (1994) Monoclonal antibodies that define canine homologues of human CD antigens: summary of the First International Canine Leukocyte Antigen Workshop (CLAW). <u>Tissue Antigens. 43 (3): 137-54.</u> Gorman, S.D. <i>et al.</i> (1994) Isolation and expression of cDNA encoding the canine CD4 and CD8 alpha antigens. <u>Tissue Antigens. 43 (3): 184-8.</u> Watson, C.J. <i>et al.</i> (1993) CD4 and CD8 monoclonal antibody therapy: strategies to prolong renal allograft survival in the dog. <u>Br J Surg. 80 (11): 1389-92.</u> Papadogiannakis, E.I. <i>et al.</i> (2009) Determination of intracellular cytokines IFN-gamma and IL-4 in canine T lymphocytes by flow cytometry following whole-blood culture. <u>Can J Vet Res. 73 (2): 137-43.</u> Benyacoub, J. <i>et al.</i> (2003) Supplementation of food with <i>Enterococcus faecium</i> (SF68) stimulates immune functions in young dogs. <u>J Nutr. 133: 1158-62.</u> Bird, R.C. <i>et al.</i> (2010) An autologous dendritic cell canine mammary tumor hybrid-cell fusion vaccine. <u>Cancer Immunol Immunother. 60: 87-97.</u> Bund, D. <i>et al.</i> (2010) Canine-DCs using different serum-free methods as an approach to provide an animal-model for immunotherapeutic strategies. <u>Cell Immunol. 263: 88-98.</u> 8. Estrela-Lima, A. <i>et al.</i> (2010) Immunophenotypic features of tumor infiltrating lymphocytes from mammary carcinomas in female dogs associated with prognostic factors and survival rates. <u>BMC Cancer. 10: 256.</u> Huang, Y.C. <i>et al.</i> (2008) CD5-low expression lymphocytes in canine peripheral blood show characteristics of natural killer cells. <u>J Leukoc Biol. 84: 1501-10.</u> Kornegay, J.N. <i>et al.</i> (2010) Widespread muscle expression of an AAV9 human

mini-dystrophin vector after intravenous injection in neonatal dystrophin-deficient dogs. <u>Mol Ther. 18: 1501-8.</u>

11. Pichavant, C. *et al.* (2010) Expression of dog microdystrophin in mouse and dog muscles by gene therapy. <u>Mol Ther. 18: 1002-9.</u>

12. Pinheiro, D. *et al* (2011) Phenotypic and functional characterization of a CD4(+) CD25(high) FOXP3(high) regulatory T-cell population in the dog. <u>Immunology. 132:</u> <u>111-22.</u>

13. Reis, A.B. *et al.* (2006) Phenotypic features of circulating leucocytes as immunological markers for clinical status and bone marrow parasite density in dogs naturally infected by *Leishmania chagasi*. <u>Clin Exp Immunol. 146: 303-11.</u>

14. Figueiredo, M.M. *et al.* (2014) Expression of Regulatory T Cells in Jejunum, Colon, and Cervical and Mesenteric Lymph Nodes of Dogs Naturally Infected with *Leishmania infantum*. Infect Immun. 82: 3704-12.

 Costa-Pereira, C. *et al.* (2015) One-year timeline kinetics of cytokine-mediated cellular immunity in dogs vaccinated against visceral leishmaniasis. <u>BMC Vet Res. 11 (1): 92.</u>
 Schaut, R.G. *et al.* (2016) Regulatory IgDhi B Cells Suppress T Cell Function via IL-10 and PD-L1 during Progressive Visceral Leishmaniasis. <u>J Immunol. Apr 13. pii: 1502678.</u> [Epub ahead of print]

17. Tagawa, M. *et al.* (2016) Evaluation of Costimulatory Molecules in Peripheral Blood Lymphocytes of Canine Patients with Histiocytic Sarcoma. <u>PLoS One. 11 (2): e0150030.</u>
18. Riondato, F. *et al.* (2015) Analytical and diagnostic validation of a flow cytometric strategy to quantify blood and marrow infiltration in dogs with large b-cell lymphoma. <u>Cytometry B Clin Cytom. Dec 13. [Epub ahead of print]</u>

19. Cortese, L. *et al.* (2015) An immune-modulating diet increases the regulatory T cells and reduces T helper 1 inflammatory response in Leishmaniosis affected dogs treated with standard therapy. <u>BMC Vet Res. 11: 295.</u>

20. Miller, J. *et al.* (2015) Humoral and Cellular Immune Response in Canine Hypothyroidism. <u>J Comp Pathol. 153 (1): 28-37.</u>

 Riondato, F. *et al.* (2016) Identification of a suitable internal control for fluorescence analysis on canine peripheral blood samples. <u>Vet Immunol Immunopathol. 172: 38-42.</u>
 Martini, V. *et al.* (2015) Canine small clear cell/T-zone lymphoma: clinical presentation and outcome in a retrospective case series. <u>Vet Comp Oncol. Jun 3. [Epub ahead of print]</u>
 Gelain, M.E. *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. <u>Vet Immunol Immunopathol. 159 (1-2): 91-6.</u>

24. Duz AL *et al.* (2014) The Tcl and Tcll *Trypanosoma cruzi* experimental infections induce distinct immune responses and cardiac fibrosis in dogs. <u>Mem Inst Oswaldo Cruz.</u> 109 (8): 1005-13.

25. Munhoz, T.D. *et al.* (2016) Regulatory T cells in dogs with multicentric lymphoma: peripheral blood quantification at diagnosis and after initial stage chemotherapy <u>Arg. Bras.</u> <u>Med. Vet. Zootec., v.68, n.1, p.1-9, 2016</u>

26. Bonnefont-Rebeix, C. *et al.* (2016) Characterization of a novel canine T-cell line established from a spontaneously occurring aggressive T-cell lymphoma with large granular cell morphology. <u>Immunobiology. 221 (1): 12-22.</u>

27. Viana, K.F. *et al.* (2015) Setting the proportion of CD4+ and CD8+ T-cells co-cultured with canine macrophages infected with *Leishmania chagasi*. <u>Vet Parasitol. 211 (3-4):</u> 124-32.

28. Bromberek, J.L. et al. (2016) Breed Distribution and Clinical Characteristics of B Cell

Chronic Lymphocytic Leukemia in Dogs. J Vet Intern Med. 30 (1): 215-22.

29. Mie K *et al.* (2016) Influence of transfusion of lymphokine-activated T killer cells on inflammatory responses in dogs after laparotomy. <u>J Vet Med Sci. Jan 2. [Epub ahead of print]</u>

30. Miglio, A. *et al.* (2014) Acute undifferentiated leukaemia in a dog. <u>Aust Vet J. 92 (12):</u> <u>499-503.</u>

31. Villaescusa, A. *et al.* (2015) Effects of doxycycline on haematology, blood chemistry and peripheral blood lymphocyte subsets of healthy dogs and dogs naturally infected with *Ehrlichia canis*. <u>Vet J. 204 (3): 263-8.</u>

32. Fiuza JA *et al.* (2015) Vaccination using live attenuated *Leishmania donovani* centrin deleted parasites induces protection in dogs against *Leishmania infantum*. <u>Vaccine. 33 (2):</u> <u>280-8.</u>

33. Perosso, J. *et al.* (2014) Alteration of sFAS and sFAS ligand expression during canine visceral leishmaniosis. <u>Vet Parasitol. 205 (3-4): 417-23.</u>

34. Heinrich, F. *et al.* (2015) Immunophenotyping of immune cell populations in the raccoon (*Procyon lotor*). <u>Vet Immunol Immunopathol. 168 (3-4): 140-6.</u>

35. Poggi A *et al.* (2016) Prognostic significance of Ki67 evaluated by flow cytometry in dogs with high-grade B-cell lymphoma. <u>Vet Comp Oncol. Jan 21. [Epub ahead of print]</u>
36. McGill, J.L. *et al.* (2016) Vaccination with an Attenuated Mutant of Ehrlichia chaffeensis Induces Pathogen-Specific CD4+ T Cell Immunity and Protection from Tick-Transmitted Wild-Type Challenge in the Canine Host. PLoS One. 11 (2): e0148229.

37. Villaescusa, A. *et al.* (2012) Evaluation of peripheral blood lymphocyte subsets in family-owned dogs naturally infected by *Ehrlichia canis*. <u>Comp Immunol Microbiol Infect</u> Dis. 35 (4): 391-6.

38. Schaut RG *et al.* (2016) Recovery of antigen-specific T cell responses from dogs infected with *Leishmania (L.) infantum* by use of vaccine associated TLR-agonist adjuvant. <u>Vaccine. 34 (44): 5225-5234</u>.

39. Miranda, S. *et al.* (2007) Characterization of circulating lymphocyte subpopulations in canine leishmaniasis throughout treatment with antimonials and allopurinol. <u>Vet Parasitol.</u> <u>144 (3-4): 251-60.</u>

40. Viana, K.F. *et al.* (2016) Application of rapid in vitro co-culture system of macrophages and T-cell subsets to assess the immunogenicity of dogs vaccinated with live attenuated *Leishmania donovani* centrin deleted parasites (LdCen-/-). <u>Parasit Vectors. 9: 250.</u>

41. Michael, H.T. *et al.* (2013) Isolation and characterization of canine natural killer cells. <u>Vet Immunol Immunopathol. 155 (3): 211-7.</u>

42. Mitchell, L. *et al.* (2012) Induction of remission results in spontaneous enhancement of anti-tumor cytotoxic T-lymphocyte activity in dogs with B cell lymphoma. <u>Vet Immunol Immunopathol. 145 (3-4): 597-603.</u>

43. DaSilva, A.V.A. *et al.* (2018) Morphophysiological changes in the splenic extracellular matrix of *Leishmania infantum*-naturally infected dogs is associated with alterations in lymphoid niches and the CD4+ T cell frequency in spleens. <u>PLoS Negl Trop Dis. 12 (4):</u> <u>e0006445.</u>

44. Roatt, B.M.*et al.* (2017) A Vaccine Therapy for Canine Visceral Leishmaniasis Promoted Significant Improvement of Clinical and Immune Status with Reduction in Parasite Burden. <u>Front Immunol. 8: 217.</u>

45. Aricò, A. *et al.* (2013) The role of vascular endothelial growth factor and matrix metalloproteinases in canine lymphoma: *in vivo* and *in vitro* study. <u>BMC Vet Res. 9: 94.</u>

NEGATIVE C Tel: +1 800 265 Fax: +1 919 878 Email: antibody_	CONTROL:Pacific Blue® (M 7376 Worldwide 3751 sales_us@bio-rad.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-		Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.con bio-rad-antibodies.com/datashee
NEGATIVE C Tel: +1 800 265 Fax: +1 919 878	ONTROL:Pacific Blue® (M 7376 Worldwide 13751	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	-	<u>1CA6004PB)</u>		
mended Ne	guire controlo			
	gative Controls			
d Produc	ts			
ory	For research purposes	s only		
nd Safety ion	•			at:
edgements	Corporation. The trans product solely in resea and the buyer must no diagnostic, therapeutic services, or information or quality assurance of research. For information as described above, or	afer of this product is cor arch, excluding contract of sell or otherwise trans of prophylactic purpose n in return for compensa r quality control, or (d) re- tion on purchasing a lice ontact Life Technologies	ntingent on the research or a fer this produces; (b) testing ation on a per esale, whethe nse to this pr Corporation	he buyer using the purchased any fee for service research, loct or its components for (a) analysis or screening r-test basis; (c) manufacturing er or not resold for use in roduct for purposes other than
90	12 months from date c	of despatch		
			-	
	-		ded. This pro	oduct is photosensitive and
	This product should be	e stored undiluted.		
	Store at +4°C or at -20	0°C if preferred.		
	Leishmune [®] , and Leis	h-Tec [®] Vaccines agains		
	edgements nd Safety on ry d Produc	Leishmune [®] , and Leis (Basel). 8 (4)Nov 17 [f Store at +4°C or at -20 This product should be Storage in frost-free fr should be protected freezin product contain a prece edgements This product is provide Corporation. The trans product solely in resea and the buyer must no diagnostic, therapeutic services, or informatio or quality assurance o research. For informatio as described above, c CA 92008 USA or out Material Safety Datast https://www.bio-rad-ar 10041 ry For research purposes	Leishmune®, and Leish-Tec® Vaccines against (Basel). 8 (4)Nov 17 [Epub ahead of print].Store at +4°C or at -20°C if preferred.This product should be stored undiluted.Storage in frost-free freezers is not recommen should be protected from light.Avoid repeated freezing and thawing as this m product contain a precipitate we recommend nre12 months from date of despatchedgementsThis product is provided under an intellectual p Corporation. The transfer of this product is cor product solely in research, excluding contract p and the buyer must not sell or otherwise transf diagnostic, therapeutic or prophylactic purpose services, or information in return for compensa or quality assurance or quality control, or (d) re research. For information on purchasing a lice as described above, contact Life Technologies CA 92008 USA or outlicensing@thermofisher.comnd Safety onMaterial Safety Datasheet documentation #100 https://www.bio-rad-antibodies.com/SDS/MCA1 10041ryFor research purposes only	Store at +4°C or at -20°C if preferred.This product should be stored undiluted.Storage in frost-free freezers is not recommended. This proshould be protected from light.Avoid repeated freezing and thawing as this may denature product contain a precipitate we recommend microcentrifugre12 months from date of despatchedgementsThis product is provided under an intellectual property licent Corporation. The transfer of this product is contingent on the product solely in research, excluding contract research or a and the buyer must not sell or otherwise transfer this product diagnostic, therapeutic or prophylactic purposes; (b) testing services, or information in return for compensation on a per or quality assurance or quality control, or (d) resale, whether research. For information on purchasing a license to this pr as described above, contact Life Technologies Corporation CA 92008 USA or outlicensing@thermofisher.comnd Safety onMaterial Safety Datasheet documentation #10041 available https://www.bio-rad-antibodies.com/SDS/MCA1039PB 10041ryFor research purposes only

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint