

Datasheet: MCA1044GA

## **BATCH NUMBER 161486**

Description:	RAT ANTI DOG MHC CLASS II MONOMORPHIC
Specificity:	MHC CLASS II MONOMORPHIC
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	YKIX334.2
Isotype:	lgG2a
Quantity:	0.1 mg

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/50 - 1/100
Immunohistology - Frozen	•			
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation	•			
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Dog
Species Cross Reactivity	Does not react with:Hooded Seal
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide

## Stabilicare

Stabilisers	
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Con A activated canine peripheral blood cells.
RRID	AB_11203404
Fusion Partners	Spleen cells from immunised DA rats were fused with cells of the rat Y3/Ag1.2.3 myeloma cell line.
Specificity	Rat anti Dog MHC Class II Monomorphic antibody, clone YKIX334.2 recognizes a monomorphic epitope on canine MHC Class II and was classified at the First Canine Leucocyte Antigen Workshop (Cobbold et al. 1994). The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In dogs, this is referred to as the dog leukocyte antigen (DLA) region. Rat anti Dog MHC Class II immunoprecipitates an antigen of ~32/34 kDa and blocks the proliferation of MHC Class II dependent responses <i>in vitro</i> . In dogs, MHC Class II is expressed by all peripheral blood mononuclear cells.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Cobbold, S. &amp; Metcalfe, S. (1994) Monoclonal antibodies that define canine homologues of human CD antigens: summary of the First International Canine Leukocyte Antigen Workshop (CLAW). <u>Tissue Antigens. 43 (3): 137-54.</u></li> <li>Watson, C.J. <i>et al.</i> (1994) Immunosuppression of canine renal allograft recipients by CD4 and CD8 monoclonal antibodies. <u>Tissue Antigens. 43 (3): 155-62.</u></li> <li>Reis, A.B. <i>et al.</i> (2006) Phenotypic features of circulating leucocytes as immunological markers for clinical status and bone marrow parasite density in dogs naturally infected by <i>Leishmania chagasi</i>. <u>Clin Exp Immunol.146: 303-11.</u></li> <li>Araújo, M.S. <i>et al.</i> (2011) Immunological changes in canine peripheral blood leukocytes triggered by immunization with first or second generation vaccines against canine visceral leishmaniasis. <u>Vet Immunol Immunopathol. 141: 64-75.</u></li> </ol>

- 5. Bonnefont-Rebeix, C. et al. (2007) Toll-like receptor 3 (TLR3): a new marker of canine monocytes-derived dendritic cells (cMo-DC). Vet Immunol Immunopathol. 2007 Jul 15;118(1-2):134-9.
- 6. Bund, D. et al. (2010) Canine-DCs using different serum-free methods as an approach to provide an animal-model for immunotherapeutic strategies. Cell Immunol. 263: 88-98.
- 7. Mito, K. et al. (2010) IFNy markedly cooperates with intratumoral dendritic cell vaccine in dog tumor models. Cancer Res. 70: 7093-101.
- 8. Sanchez, M.A. et al. (2004) Organ-specific immunity in canine visceral leishmaniasis: analysis of symptomatic and asymptomatic dogs naturally infected with Leishmania chagasi. Am J Trop Med Hyg. 70: 618-24.
- 9. Schütze, N. et al. (2009) Inactivated parapoxvirus ovis activates canine blood phagocytes and T lymphocytes. Vet Microbiol. 137: 260-7.

- 10. Liu, Y. *et al.* (2000) Immunosuppressant-free allotransplantation of the tracheaThe antigenicity of tracheal grafts can be reduced by removing the epithelium and mixed glands from the graft by detergent treatment. <u>J Thorac Cardiovasc Surg. 120: 108-14.</u>

  11. Larsen, A.K. *et al.* (2013) Entry and elimination of marine mammal Brucella spp. by hooded seal (Cystophora cristata) alveolar macrophages *in vitro*. <u>PLoS One. 8: e70186.</u>

  12. Bonnefont-Rebeix, C. *et al.* (2016) Characterization of a novel canine T-cell line established from a spontaneously occurring aggressive T-cell lymphoma with large granular cell morphology. <u>Immunobiology. 221 (1): 12-22.</u>
- 13. Lin, S-C. *et al.* (2014) Immune Characterization of Peripheral Blood Mononuclear cells of the Dogs Restored from Innoculation of Canine Transmissible Venereal Tumor Cells. Tai Vet J. 40 (04): 181-90.
- 14. Constantinoiu, C.C. *et al.* (2015) Mucosal tolerance of the hookworm Ancylostoma caninum in the gut of naturally infected wild dogs. <u>Parasite Immunol. 37 (10): 510-20.</u>
  15. Lu, T. *et al.* (2017) Effects of cryopreservation on tracheal allograft antigenicity in dogs. J Thorac Dis. 9 (7): 2038-2047.
- 16. Reineking, W. *et al.* (2018) Canine primary jejunal and colonic epithelial cells predominantly express TLR5 and TLR9 but do not change TLR expression pattern after stimulation with certain Toll-like receptor ligands. <u>Vet Immunol Immunopathol. 206: 16-24.</u>
  17. Martini, V. *et al.* (2019) Prognostic role of non-neoplastic lymphocytes in lymph node aspirates from dogs with diffuse large B-cell lymphoma treated with chemo-immunotherapy. Res Vet Sci. 125: 130-5.
- 18. Grudzien, M. *et al.* (2021) A newly established canine NK-type cell line and its cytotoxic properties. Vet Comp Oncol. 19 (3): 567-77.
- 19. Bragato, J.P. *et al.* (2022) miRNA-21 regulates CD69 and IL-10 expression in canine leishmaniasis. PLoS One. 17 (3): e0265192.

#### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1044GA">https://www.bio-rad-antibodies.com/SDS/MCA1044GA</a> 10040
Regulatory	For research purposes only

## Related Products

## **Recommended Secondary Antibodies**

Rabbit Anti Rat IgG (STAR16...) <u>DyLight®800</u>

Rabbit Anti Rat IgG (STAR17...) FITC

Goat Anti Rat IgG (STAR73...) RPE

Rabbit Anti Rat IgG (STAR21...) HRP

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) DyLight®550, DyLight®650, DyLight®800

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

Goat Anti Rat IgG (STAR72...)

HRP
Goat Anti Rat IgG (STAR69...)

FITC

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

 $\textbf{Email: antibody\_sales\_us@bio-rad.com} \\ \textbf{Email: antibody\_sales\_uk@bio-rad.com} \\ \textbf{Email: antibody\_sales\_uk@b$ 

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M382476:210513'

## Printed on 28 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint