

Datasheet: MCA1053F BATCH NUMBER 150242

Description:	MOUSE ANTI HUMAN CD48:FITC
Specificity:	CD48
Other names:	BLAST-1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MEM-102
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications

Buffer Solution

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross	Reacts with: Rhesus Monkey
Reactivity	N.B. Antibody reactivity and working conditions may vary bety

Phosphate buffered saline

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG conju	ugated to Fluorescein Isoth	niocyanate Isomer 1
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prep	ared by affinity chromatog	raphy on Protein A

Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
External Database Links	UniProt: P09326 Related reagents		
	Entrez Gene: 962 CD48 Related reagents		
Synonyms	BCM1, BLAST1		
RRID	AB_321437		
Specificity	Mouse anti Human CD48 antibody, clone MEM-102 recognizes human CD48, also known as B-lymphocyte activation marker BLAST-1, BCM1 surface antigen, Leukocyte antigen MEM-102, SLAM family member 2 or Signaling lymphocytic activation molecule 2. CD 48, as a mature molecule is a 193 amino acid, ~45 kDa cell membrane glycoprotein containing 2 lg-like domains.		
	CD48 acts as a counter-receptor for CD2 (<u>Kato et al. 1992</u>) and is involved in immune regulation and tolerance (<u>McArdel et al. 2016</u>)		
	Mouse anti Human CD48 antibody, clone MEM-102 is useful for the visualisation of CD48 expressing cells by immunofluorescence (Enose-Akahata et al. 2009) and flow cytometry and may be used in studies to differentiate haemapoietic tumors, sarcomas and melanomas.		
Flow Cytometry	Use 10ul of the suggested working dilution to label 1 x 10^6 cells in 100ul.		
References	1. Horejsí, V. <i>et al.</i> (1988) Monoclonal antibodies against human leucocyte antigens. II. Antibodies against CD45 (T200), CD3 (T3), CD43, CD10 (CALLA), transferrin receptor (T9), a novel broadly expressed 18-kDa antigen (MEM-43) and a novel antigen of restricted expression (MEM-74). Folia Biol (Praha). 34 (1): 23-34. 2. Korínek, V. <i>et al.</i> (1991) The human leucocyte antigen CD48 (MEM-102) is closely related to the activation marker Blast-1. Immunogenetics. 33 (2): 108-12. 3. Cinek, T. & Horejsí, V. (1992) The nature of large noncovalent complexes containing glycosyl-phosphatidylinositol-anchored membrane glycoproteins and protein tyrosine kinases. J Immunol. 149 (7): 2262-70. 4. Staffler, G. <i>et al.</i> (2003) Selective inhibition of T cell activation via CD147 through novel modulation of lipid rafts. J Immunol. 171 (4): 1707-14.		

6. Enose-Akahata, Y. *et al.* (2009) High expression of CD244 and SAP regulated CD8 T cell responses of patients with HTLV-I associated neurologic disease. <u>PLoS Pathog. 5:</u>

e1000682.

- 7. Fukushima, K. *et al.* (2005) Functional role played by the glycosylphosphatidylinositol anchor glycan of CD48 in interleukin-18-induced interferon-gamma production. <u>J Biol Chem.</u> 280: 18056-62.
- 8. Kanuga, N. *et al.* (2002) Characterization of genetically modified human retinal pigment epithelial cells developed for in vitro and transplantation studies. <u>Invest Ophthalmol Vis Sci. 43: 546-55.</u>
- 9. Peterson, L.A. *et al.* (2019) Individual differences in the response of human β-lymphoblastoid cells to the cytotoxic, mutagenic and DNA damaging effects of a DNA methylating agent, *N*-methylnitrosourethane. <u>Chem Res Toxicol</u>. <u>Oct 07 [Epub ahead of print]</u>.

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1053F 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

 ${\bf Email: antibody_sales_de@bio-rad.com}$

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M364716:200529'

Printed on 18 Jan 2024