

## Datasheet: MCA1054A488

Description:	MOUSE ANTI HUMAN CD59:Alexa Fluor® 488
Specificity:	CD59
Other names:	HRF, PROTECTIN
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	MEM-43
Isotype:	lgG2a
Quantity:	100 TESTS/1ml

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes N	10	Not Determined	Suggested Dilution
	Flow Cytometry	-			Neat
	Where this antibody h	as not been test	ed for u	ise in a particular teo	chnique this does not
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.				
Target Species	Human				
Product Form	Purified IgG conjugated to Alexa Fluor 488 - liquid				
Max Ex/Em	Fluorophore	Excitation Max	(nm) I	Emission Max (nm)	
	Alexa Fluor®488	495		519	
Preparation	Purified IgG prepared supernatant	by affinity chrom	natogra	phy on Protein A froi	m tissue culture
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% sodium azide (NaN <sub>3</sub> ) 1% bovine serum albumin				
Stabilisers					
Approx. Protein Concentrations	IgG concentration 0.0	5 mg/ml			

Immunogen	Thymocytes and T lymphocytes.
External Database Links	UniProt: <u>P13987</u> <u>Related reagents</u> Entrez Gene: <u>966</u> CD59 <u>Related reagents</u>
Synonyms	MIC11, MIN1, MIN2, MIN3, MSK21
RRID	AB_2076229
Specificity	<b>Mouse anti Human CD59 antibody, clone MEM-43</b> recognizes CD59, a glycosyl- phosphatidylinositol (GPI) anchored membrane protein also known as membrane attack complex inhibition factor. CD59 blocks the formation of the complement membrane attack complex (MAC) by binding of C8a and C9. CD59 is found on all types of leucocytes including platelets and is also expressed on many non-haematopoietic cells. The epitope recognized by Mouse anti Human CD59 antibody, clone MEM-43 is lost after reduction therefore, non-reducing conditions are required for western blotting techniques.
Flow Cytometry	Use 10µl of the suggested working dilution to label $10^6$ cells or $100µl$ whole blood
References	<ol> <li>Horejsí, V. <i>et al.</i> (1988) Monoclonal antibodies against human leucocyte antigens. II. Antibodies against CD45 (T200), CD3 (T3), CD43, CD10 (CALLA), transferrin receptor (T9), a novel broadly expressed 18-kDa antigen (MEM-43) and a novel antigen of restricted expression (MEM-74). Folia Biol (Praha). 34 (1): 23-34.</li> <li>Stefanová, I. <i>et al.</i> (1989) Characterization of a broadly expressed human leucocyte surface antigen MEM-43 anchored in membrane through phosphatidylinositol. Mol Immunol. 26 (2): 153-61.</li> <li>Stefanová, I. <i>et al.</i> (1989) in Leucocyte Typing IV: White cell differentiation antigens. Ed. Knapp, W. <i>et al.</i> Oxford University Press pp 678-97.</li> <li>Stefanová, I. &amp; Horejsí, V. (1991) Association of the CD59 and CD55 cell surface glycoproteins with other membrane molecules. J Immunol. 147 (5): 1587-92.</li> <li>Tandon, N. <i>et al.</i> (1994) Expression and function of multiple regulators of complement activation in autoimmune thyroid disease. Immunology. 81 (4): 643-7.</li> <li>Vanderplasschen, A. <i>et al.</i> (1997) Extracellular enveloped vaccinia virus is resistant to complement because of incorporation of host complement control proteins into its envelope. <u>Proc Natl Acad Sci U S A. 95: 7544-9</u>.</li> <li>Cowan, P.J. <i>et al.</i> (1998) High-level endothelial expression of human CD59 prolongs heart function in an <i>ex vivo</i> model of xenograft rejection. <u>Transplantation. 65: 826-31.</u></li> <li>Chong, Y.H. and Lee, M.J. (2000) Expression of complement inhibitor protein CD59 in human neuronal and glial cell lines treated with HIV-1 gp41 peptides. <u>J Neurovirol. 6:</u> <u>51-60.</u></li> <li>Shamri, R. <i>et al.</i> (2002) Chemokine stimulation of lymphocyte alpha 4 integrin avidity but not of leukocyte function-associated antigen-1 avidity to endothelial ligands under shear flow requires cholesterol membrane rafts. J Biol Chem. 277: 40027-35.</li> <li>Zhang, J. <i>et al.</i> (2002) Early complement activation and decreased levels of</li> </ol>

	glycosylphosphatidylinositol-anchored complement inhibitors in human and experimental
	diabetic retinopathy. <u>Diabetes. 51: 3499-504.</u>
	11. Donin, N. et al. (2003) Complement resistance of human carcinoma cells depends on
	membrane regulatory proteins, protein kinases and sialic acid. <u>Clin Exp Immunol. 131:</u> 254-63.
	12. Gendek-Kubiak, H. and Gendek, E.G. (2004) Immunolocalization of protectin (CD59)
	and macrophages in polymyositis and dermatomyositis. <u>J Neuroimmunol. 149: 187-94.</u>
	13. Jolly, C, and Sattentau. Q.J. (2005) Human Immunodeficiency Virus Type 1 Virological
	Synapse Formation in T Cells Requires Lipid Raft Integrity <u>J Virol. 79: 12088-94.</u>
	14. Ohyama, M. <i>et al.</i> (2006) Characterization and isolation of stem cell-enriched human
	hair follicle bulge cells. <u>J Clin Invest. 116: 249-60.</u>
	15. Ellison, B.S. <i>et al.</i> (2007) Complement susceptibility in glutamine deprived breast
	cancer cells. <u>Cell Div. 2007 2: 20.</u>
	16. Takemoto, M. <i>et al.</i> (2007) Human herpesvirus 7 infection increases the expression
	levels of CD46 and CD59 in target cells. <u>J Gen Virol. 88: 1415-22.</u>
	17. Shaw, M.L. <i>et al.</i> (2008) Cellular proteins in influenza virus particles. <u>PLoS Pathog. 4:</u>
	e1000085.
	18. Bonnon, C. <i>et al.</i> (2010) Selective export of human GPI-anchored proteins from the
	endoplasmic reticulum. <u>J Cell Sci. 123: 1705-15.</u>
	19. Sadallah, S. <i>et al.</i> (2011) Microparticles (ectosomes) shed by stored human platelets
	downregulate macrophages and modify the development of dendritic cells. <u>J Immunol.</u>
	186: 6543-52.
	20. Rondelli, T. <i>et al.</i> (2013) The frequency of granulocytes with spontaneous somatic
	mutations: a wide distribution in a normal human population. PLoS One. 8 (1): e54046.
	21. Abe, Y. et al. (2017) Glycan region of GPI anchored-protein is required for cytocidal
	oligomerization of an anticancer parasporin-2, Cry46Aa1 protein, from Bacillus
	thuringiensis strain A1547. J Invertebr Pathol. 142: 71-81.
	22. Sica, M. et al. (2017) Eculizumab treatment: stochastic occurrence of C3 binding to
	individual PNH erythrocytes. <u>J Hematol Oncol. 10 (1): 126.</u>
	23. Gullipalli, D. et al. (2018) Antibody Inhibition of Properdin Prevents Complement-
	Mediated Intravascular and Extravascular Hemolysis. <u>J Immunol. 201 (3): 1021-1029.</u>
	24. Ueda, M. et al. (2019) Endovascular trophoblast expresses CD59 to evade
	complement-dependent cytotoxicity. Mol Cell Endocrinol. 490: 57-67.
<u> </u>	
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at 2000 on received when the word aligned the approach of 2000 for
	-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for
	short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in
	frost-free freezers is not recommended. This product is photosensitive and should be
	protected from light.
Guarantee	12 months from date of despatch
	·
Acknowledgements	This product is provided under an intellectual property licence from Life Technologies
	Corporation. The transfer of this product is contingent on the buyer using the purchase
	product solely in research, excluding contract research or any fee for service research,
	and the buyer must not sell or otherwise transfer this product or its components for (a)

	diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in
	research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1054A488 10041
Regulatory	For research purposes only
Related Produ	cts
Recommended N	egative Controls

MOUSE IgG2a NEGATIVE CONTROL: Alexa Fluor® 488 (MCA929A488)

## **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-r	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M410477:221028'

## Printed on 26 Jun 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint