

Datasheet: MCA1054GA BATCH NUMBER 0515

Description:	MOUSE ANTI HUMAN CD59		
Specificity:	CD59		
Other names:	HRF, PROTECTIN		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	MEM-43		
Isotype:	lgG2a		
Quantity:	0.1 mg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	•			1/5 - 1/10
	Immunohistology - Frozen	-			
	Immunohistology - Paraffin (1)	-			
	ELISA	-			
	Immunoprecipitation	-			
	Western Blotting (2)	-			
	Immunofluorescence	-			
	Immuno-electron Microscopy				
Where this antibody has not been tested for use in a particular technique necessarily exclude its use in such procedures. Suggested working dilut a guide only. It is recommended that the user titrates the antibody for us system using appropriate negative/positive controls. (1) This product requires antigen retrieval using heat treatment prio paraffin sections.Sodium citrate buffer pH 6.0 is recommended for (2) This product recognizes CD59 under non-reducing conditions.				g dilutions are given a for use in their own nt prior to staining of ed for this purpose.	
Farget Species	Human				
Product Form	Purified IgG - liquid				

Preparation	Purified IgG prepared by affinity chromatography on Protein A			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide			
Approx. Protein Concentrations	IgG concentration 1 mg/ml			
Immunogen	Thymocytes and T lymphocytes.			
External Database Links	UniProt: <u>P13987</u> <u>Related reagents</u> Entrez Gene:			
	966 CD59 Related reagents			
Synonyms	MIC11, MIN1, MIN2, MIN3, MSK21			
RRID	AB_323963			
Specificity	Mouse anti Human CD59 antibody, clone MEM-43 recognizes CD59, a glycosyl- phosphatidylinositol (GPI) anchored membrane protein also known as membrane attack complex inhibition factor. CD59 blocks the formation of the complement membrane attack complex (MAC) by binding of C8a and C9. CD59 is found on all types of leucocytes including platelets and is also expressed on many non-haematopoietic cells. The epitope recognized by Mouse anti Human CD59 antibody, clone MEM-43 is lost after reduction therefore, non-reducing conditions are required for western blotting techniques.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.			
Histology Positive Control Tissue	Tonsil			
References	 Stefanova, I. <i>et al.</i> (1989) in Leucocyte Typing IV: White cell differentiation antigens. Ed. Knapp, W. <i>et al.</i> Oxford University Press pp 678-97. Stefanová, I. <i>et al.</i> (1989) Characterization of a broadly expressed human leucocyte surface antigen MEM-43 anchored in membrane through phosphatidylinositol. <u>Mol</u> <u>Immunol. 26 (2): 153-61.</u> Tandon, N. <i>et al.</i> (1994) Expression and function of multiple regulators of complement activation in autoimmune thyroid disease. <u>Immunology. 81 (4): 643-7.</u> Horejsí, V. <i>et al.</i> (1988) Monoclonal antibodies against human leucocyte antigens. II. Antibodies against CD45 (T200), CD3 (T3), CD43, CD10 (CALLA), transferrin receptor (T9), a novel broadly expressed 18-kDa antigen (MEM-43) and a novel antigen of restricted expression (MEM-74). <u>Folia Biol (Praha). 34 (1): 23-34.</u> 			

6. Shaw, M.L. *et al.* (2008) Cellular proteins in influenza virus particles. <u>PLoS Pathog. 4:</u> <u>e1000085.</u>

7. Sadallah, S. *et al.* (2011) Microparticles (ectosomes) shed by stored human platelets downregulate macrophages and modify the development of dendritic cells. <u>J Immunol.</u> <u>186: 6543-52.</u>

8. Jolly, C, and Sattentau. Q.J. (2005) Human Immunodeficiency Virus Type 1 Virological Synapse Formation in T Cells Requires Lipid Raft Integrity <u>J Virol. 79: 12088-94.</u>

9. Shamri, R. *et al.* (2002) Chemokine stimulation of lymphocyte alpha 4 integrin avidity but not of leukocyte function-associated antigen-1 avidity to endothelial ligands under shear flow requires cholesterol membrane rafts. <u>J Biol Chem. 277: 40027-35.</u>

10. Bonnon, C. *et al.* (2010) Selective export of human GPI-anchored proteins from the endoplasmic reticulum. <u>J Cell Sci. 123: 1705-15.</u>

11. Zhang, J. *et al.* (2002) Early complement activation and decreased levels of glycosylphosphatidylinositol-anchored complement inhibitors in human and experimental diabetic retinopathy. <u>Diabetes. 51: 3499-504.</u>

12. Ellison, B.S. *et al.* (2007) Complement susceptibility in glutamine deprived breast cancer cells. <u>Cell Div. 2007 2: 20.</u>

 Cowan, P.J. *et al.* (1998) High-level endothelial expression of human CD59 prolongs heart function in an *ex vivo* model of xenograft rejection. <u>Transplantation. 65: 826-31.</u>
 Vanderplasschen, A. *et al.* (1997) Extracellular enveloped vaccinia virus is resistant to complement because of incorporation of host complement control proteins into its envelope. Proc Natl Acad Sci U S A. 95: 7544-9.

15. Takemoto, M. *et al.* (2007) Human herpesvirus 7 infection increases the expression levels of CD46 and CD59 in target cells. J Gen Virol. 88: 1415-22.

16. Chong, Y.H. and Lee, M.J. (2000) Expression of complement inhibitor protein CD59 in human neuronal and glial cell lines treated with HIV-1 gp41 peptides. J Neurovirol. 6: 51-60.

Gendek-Kubiak, H. and Gendek, E.G. (2004) Immunolocalization of protectin (CD59) and macrophages in polymyositis and dermatomyositis. <u>J Neuroimmunol. 149: 187-94.</u>
 Abe, Y. *et al.* (2017) Glycan region of GPI anchored-protein is required for cytocidal oligomerization of an anticancer parasporin-2, Cry46Aa1 protein, from *Bacillus thuringiensis* strain A1547. J Invertebr Pathol. 142: 71-81.

19. Sica, M. *et al.* (2017) Eculizumab treatment: stochastic occurrence of C3 binding to individual PNH erythrocytes. J Hematol Oncol. 10 (1): 126.

20. Ueda, M. *et al.* (2019) Endovascular trophoblast expresses CD59 to evade complement-dependent cytotoxicity. <u>Mol Cell Endocrinol. 490: 57-67.</u>

21. Rondelli, T. *et al.* (2013) The frequency of granulocytes with spontaneous somatic mutations: a wide distribution in a normal human population. <u>PLoS One. 8 (1): e54046.</u>

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch			
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1054GA 10040			
Regulatory	For research purposes only			
Related Produc	cts			
Recommended Se	econdary Antibodies			
Rabbit Anti Mouse IgG (STAR12) RPE				

Rabbit Anti Mouse IgG (STAR12)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	RPE
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (STAR70)	FITC
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,
	<u>DyLight®650</u> , <u>DyLight®680</u> , <u>DyLight®800</u> ,
	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	FITC
Goat Anti Mouse IgG (STAR77)	HRP
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP
Recommended Negative Controls	
	20)

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50	
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M364726:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint