

Datasheet: MCA1054XZ

Description:	MOUSE ANTI HUMAN CD59:Preservative Free	
Specificity:	CD59	
Other names:	HRF, PROTECTIN	
Format:	Preservative Free	
Product Type:	Monoclonal Antibody	
Clone:	MEM-43	
Isotype:	lgG2a	
Quantity:	1 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/5 - 1/10
Immunohistology - Frozen	•			
Immunohistology - Paraffin (1)	•			
ELISA				
Immunoprecipitation				
Western Blotting (2)	•			
Immunofluorescence	•			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose. (2)This product recognizes CD59 under non-reducing conditions.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	A from tissue culture

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Thymocytes and T lymphocytes.
External Database Links	UniProt: P13987 Related reagents Entrez Gene: 966 CD59 Related reagents
Synonyms	MIC11, MIN1, MIN2, MIN3, MSK21
RRID	AB_323835
Specificity	Mouse anti Human CD59 antibody, clone MEM-43 recognizes CD59, a glycosylphosphatidylinositol (GPI) anchored membrane protein also known as membrane attack complex inhibition factor. CD59 blocks the formation of the complement membrane attack complex (MAC) by binding of C8a and C9. CD59 is found on all types of leucocytes including platelets and is also expressed on many non-haematopoietic cells. The epitope recognized by Mouse anti Human CD59 antibody, clone MEM-43 is lost after reduction therefore, non-reducing conditions are required for western blotting techniques.
Flow Cytometry	Use 10μl of the suggested working dilution to label 10 ⁶ cells in 100μl
Histology Positive Control Tissue	Tonsil
References	1. Horejsí, V. et al. (1988) Monoclonal antibodies against human leucocyte antigens. II. Antibodies against CD45 (T200), CD3 (T3), CD43, CD10 (CALLA), transferrin receptor (T9), a novel broadly expressed 18-kDa antigen (MEM-43) and a novel antigen of restricted expression (MEM-74). Folia Biol (Praha). 34 (1): 23-34. 2. Stefanová, I. et al. (1989) Characterization of a broadly expressed human leucocyte surface antigen MEM-43 anchored in membrane through phosphatidylinositol. Mol Immunol. 26 (2): 153-61. 3. Stefanová, I. et al. (1989) in Leucocyte Typing IV: White cell differentiation antigens. Ed Knapp, W. et al. Oxford University Press pp 678-97. 4. Stefanová, I. & Horejsí, V. (1991) Association of the CD59 and CD55 cell surface glycoproteins with other membrane molecules. J Immunol. 147 (5): 1587-92. 5. Tandon, N. et al. (1994) Expression and function of multiple regulators of complement activation in autoimmune thyroid disease. Immunology. 81 (4): 643-7. 6. Vanderplasschen, A. et al. (1997) Extracellular enveloped vaccinia virus is resistant to complement because of incorporation of host complement control proteins into its

envelope. Proc Natl Acad Sci U S A. 95: 7544-9.

- 7. Cowan, P.J. *et al.* (1998) High-level endothelial expression of human CD59 prolongs heart function in an *ex vivo* model of xenograft rejection. <u>Transplantation</u>. 65: 826-31.
- 8. Chong, Y.H. and Lee, M.J. (2000) Expression of complement inhibitor protein CD59 in human neuronal and glial cell lines treated with HIV-1 gp41 peptides. <u>J Neurovirol. 6:</u> 51-60.
- 9. Shamri, R. *et al.* (2002) Chemokine stimulation of lymphocyte alpha 4 integrin avidity but not of leukocyte function-associated antigen-1 avidity to endothelial ligands under shear flow requires cholesterol membrane rafts. <u>J Biol Chem. 277: 40027-35.</u>
- 10. Zhang, J. *et al.* (2002) Early complement activation and decreased levels of glycosylphosphatidylinositol-anchored complement inhibitors in human and experimental diabetic retinopathy. Diabetes. 51: 3499-504.
- 11. Donin, N. *et al.* (2003) Complement resistance of human carcinoma cells depends on membrane regulatory proteins, protein kinases and sialic acid. <u>Clin Exp Immunol. 131:</u> <u>254-63.</u>
- 12. Gendek-Kubiak, H. and Gendek, E.G. (2004) Immunolocalization of protectin (CD59) and macrophages in polymyositis and dermatomyositis. <u>J Neuroimmunol</u>. 149: 187-94.
- 13. Jolly, C, and Sattentau. Q.J. (2005) Human Immunodeficiency Virus Type 1 Virological Synapse Formation in T Cells Requires Lipid Raft Integrity <u>J Virol. 79: 12088-94.</u>
- 14. Ohyama, M. *et al.* (2006) Characterization and isolation of stem cell-enriched human hair follicle bulge cells. <u>J Clin Invest.</u> 116: 249-60.
- 15. Ellison, B.S. *et al.* (2007) Complement susceptibility in glutamine deprived breast cancer cells. Cell Div. 2007 2: 20.
- 16. Takemoto, M. *et al.* (2007) Human herpesvirus 7 infection increases the expression levels of CD46 and CD59 in target cells. J Gen Virol. 88: 1415-22.
- 17. Shaw, M.L. *et al.* (2008) Cellular proteins in influenza virus particles. <u>PLoS Pathog. 4:</u> e1000085.
- 18. Bonnon, C. *et al.* (2010) Selective export of human GPI-anchored proteins from the endoplasmic reticulum. J Cell Sci. 123: 1705-15.
- 19. Sadallah, S. *et al.* (2011) Microparticles (ectosomes) shed by stored human platelets downregulate macrophages and modify the development of dendritic cells. <u>J Immunol.</u> 186: 6543-52.
- 20. Rondelli, T. *et al.* (2013) The frequency of granulocytes with spontaneous somatic mutations: a wide distribution in a normal human population. PLoS One. 8 (1): e54046.
- 21. Abe, Y. *et al.* (2017) Glycan region of GPI anchored-protein is required for cytocidal oligomerization of an anticancer parasporin-2, Cry46Aa1 protein, from *Bacillus thuringiensis* strain A1547. <u>J Invertebr Pathol. 142: 71-81.</u>
- 22. Sica, M. *et al.* (2017) Eculizumab treatment: stochastic occurrence of C3 binding to individual PNH erythrocytes. <u>J Hematol Oncol. 10 (1): 126.</u>
- 23. Gullipalli, D. *et al.* (2018) Antibody Inhibition of Properdin Prevents Complement-Mediated Intravascular and Extravascular Hemolysis. <u>J Immunol. 201 (3): 1021-1029.</u>
 24. Ueda, M. *et al.* (2019) Endovascular trophoblast expresses CD59 to evade
- complement-dependent cytotoxicity. Mol Cell Endocrinol. 490: 57-67.

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing

as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA1054XZ 10162
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) HRP
Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP
Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR70...) FITC

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M421847:230731'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint