

Datasheet: MCA1075PE BATCH NUMBER 1603

Description:	MOUSE ANTI HUMAN CD32:RPE			
Specificity:	CD32			
Other names:	FcRII			
Format:	RPE			
Product Type:	Monoclonal Antibody			
Clone:	AT10			
lsotype:	lgG1			
Quantity:	100 TESTS			

Product Details

Applications	This product has been reported to work in the following applications. This informati derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes No	Not Determined	Suggested Dilution			
	Flow Cytometry	•		Neat			
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given a a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.						
Target Species	Human						
Species Cross Reactivity	Reacts with: Dog, Rhesus Monkey, Pig N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.						
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized						
Reconstitution	Reconstitute with 1.0 ml distilled water						
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)				
	RPE 488nm laser	496	578				
Preparation	Purified IgG prepared by affinity chromatography on Protein A						

Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azide				
Stabilisers	1% Bovine Serum Albumin				
Immunogen	K562 cell line.				
External Database					
Links	UniProt:				
	P12318 Related reagents				
	P31994 Related reagents				
	P31995 Related reagents				
	Entrez Gene:				
	2212 FCGR2A Related reagents				
	9103 FCGR2C Related reagents				
Synonyms	CD32, FCG2, FCGR2A1, IGFR2				
RRID	AB_321661				
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.				
Specificity	 Mouse anti Human CD32 antibody, clone AT10 recognizes the human CD32 antigen, a ~40 kDa glycoprotein that acts as a low affinity receptor for IgG (also known as Fc gamma RII). CD32 mediates several functions including endocytosis, activation of secretion, cytotoxicity and immunomodulation. CD32 is expressed by B cells, monocytes, granulocytes and platelets. Mouse anti Human CD32 antibody, clone AT10 blocks the binding of IgG to Fc gamma RII (Larsson <i>et al.</i> 1997). 				
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.				
References	 Van Den Herik-Oudijk, I.E. <i>et al.</i> (1994) Functional analysis of human Fc gamma RII (CD32) isoforms expressed in B lymphocytes. <u>J Immunol. 152 (2): 574-85.</u> Lilliehöök, I. <i>et al.</i> (1998) Expression of adhesion and Fcgamma-receptors on canine blood eosinophils and neutrophils studied by anti-human monoclonal antibodies. <u>Vet Immunol Immunopathol. 61 (2-4): 181-93.</u> Larsson M <i>et al.</i> (1997) Human dendritic cells handling of binding, uptake and degradation of free and IgG-immune complexed dinitrophenylated human serum albumin <i>in vitro</i>. <u>Immunology. 90 (1): 138-46.</u> Mold, C. and Du Clos, T.W. (2006) C-reactive protein increases cytokine responses to <i>Streptococcus pneumoniae</i> through interactions with Fc gamma receptors. <u>J Immunol. 176: 7598-604.</u> 				

5. Dutertre, C.A. *et al.* (2008) A novel subset of NK cells expressing high levels of inhibitory FcgammaRIIB modulating antibody-dependent function. J Leukoc Biol. 84 (6): 1511-20.

6. Devriendt, B. *et al.* (2010) Targeting of *Escherichia coli* F4 fimbriae to Fcgamma receptors enhances the maturation of porcine dendritic cells. <u>Vet Immunol Immunopathol.</u> <u>135: 188-98.</u>

7. Sims, G.P. *et al.* (2005) Identification and characterization of circulating human transitional B cells. <u>Blood. 105: 4390-8.</u>

8. Benitez-Ribas, D. *et al.* (2006) Plasmacytoid dendritic cells of melanoma patients present exogenous proteins to CD4+ T cells after Fc gamma RII-mediated uptake. <u>J Exp</u> <u>Med. 203: 1629-35.</u>

9. Zhao, X.W. *et al.* (2011) CD47-signal regulatory protein-α (SIRPα) interactions form a barrier for antibody-mediated tumor cell destruction. <u>Proc Natl Acad Sci U S A. 108 (45)</u>: 18342-7.

10. Araújo, M.S.*et al.* (2011) Immunological changes in canine peripheral blood leukocytes triggered by immunization with first or second generation vaccines against canine visceral leishmaniasis. <u>Vet Immunol Immunopathol. 141: 64-75.</u>

11. Bonnefont-Rebeix, C. *et al.* (2006) CD86 molecule is a specific marker for canine monocyte-derived dendritic cells. <u>Vet Immunol Immunopathol. 109 (1-2): 167-76.</u>

12. Santer, D.M. *et al.* (2010) C1q deficiency leads to the defective suppression of IFN-alpha in response to nucleoprotein containing immune complexes. J Immunol. 185: 4738-49.

13. Shannon, O. *et al.* (2010) Platelet activation and biofilm formation by *Aerococcus urinae*, an endocarditis-causing pathogen. <u>Infect Immun. 78: 4268-75.</u>

14. Ito, T. *et al.* (1999) A CD1a+/CD11c+ subset of human blood dendritic cells is a direct precursor of Langerhans cells. J Immunol. 163: 1409-19.

15. Moreira, M.L. *et al.* (2016) Vaccination against canine leishmaniosis increases the phagocytic activity, nitric oxide production and expression of cell activation/migration molecules in neutrophils and monocytes. <u>Vet Parasitol. 220: 33-45.</u>

16. Gazendam, R.P. *et al.* (2016) Impaired killing of *Candida albicans* by granulocytes mobilized for transfusion purposes: a role for granule components. <u>Haematologica. 101</u> (5): 587-96.

17. Liu M *et al.* (2011) Vitellogenin mediates phagocytosis through interaction with FcγR. <u>Mol Immunol. 49 (1-2): 211-8.</u>

18. Petersson, F. *et al.* (2018) Platelet activation and aggregation by the opportunistic pathogen *Cutibacterium (Propionibacterium) acnes.* <u>PLoS One. 13 (1): e0192051.</u>

19. Kahn, F. *et al.* (2008) Antibodies against a surface protein of *Streptococcus pyogenes* promote a pathological inflammatory response. <u>PLoS Pathog. 4 (9): e1000149.</u>

20. Bruggeman, C.W. *et al.* (2019) Tissue-specific expression of IgG receptors by human macrophages *ex vivo*. <u>PLoS One. 14 (10): e0223264.</u>

21. Chen, T. *et al.* (2020) Capsular glycan recognition provides antibody-mediated immunity against tuberculosis. J Clin Invest. 130 (4): 1808-22.

22. Teuben, M.P.J. *et al.* (2021) Standardized porcine unilateral femoral nailing is associated with changes in PMN activation status, rather than aberrant systemic PMN prevalence. <u>Eur J Trauma Emerg Surg. Jun 10 [Epub ahead of print].</u>

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be
protected from light. Should this product contain a precipitate we recommend
microcentrifugation before use.Guarantee12 months from date of despatchHealth And Safety
InformationMaterial Safety Datasheet documentation #20487 available at:
https://www.bio-rad-antibodies.com/SDS/MCA1075PE
20487RegulatoryFor research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375257:210104'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint