

# Datasheet: MCA1076SBB675

Description:	MOUSE ANTI HUMAN CD62L:StarBright Blue 675
Specificity:	CD62L
Other names:	LECAM-1, L-SELECTIN
Format:	StarBright Blue 675
Product Type:	Monoclonal Antibody
Clone:	FMC46
Isotype:	lgG2b
Quantity:	100 TESTS/0.5ml

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No Not I	Determined	Suggested Dilution	
	Flow Cytometry	•			Neat	
	Where this product ha necessarily exclude its a guide only. It is reco system using appropri	s use in such p mmended that	rocedures. Sug the user titrates	gested working	g dilutions are given as	
Target Species	Human					
Species Cross Reactivity	Reacts with: Bovine, Cynomolgus monkey, Rhesus Monkey, Dog <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG conjugated to StarBright Blue 675 - liquid					
Max Ex/Em	Fluorophore	Excitation Ma	k (nm) Emissio	on Max (nm)		
	StarBright Blue 675	476		675		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant					
Buffer Solution	Phosphate buffered sa	aline				

Preservative Stabilisers	0.09% sodium azide (NaN <sub>3</sub> ) 1% bovine serum albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20			
Immunogen	PHA stimulated lymphoblasts			
External Database Links	UniProt: <u>P14151</u> <u>Related reagents</u> Entrez Gene: <u>6402</u> SELL <u>Related reagents</u>			
Synonyms	LNHR, LYAM1			
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line			
Specificity	<b>Mouse anti Human CD62L antibody, clone FMC46</b> recognizes human CD62L, also known a L-selectin, a 74-95 kDa member of the selectin family of adhesion receptors, which acts as a ligand for both CD62P (P-selectin) and CD62E (E-selectin). Human CD62L is constitutively expressed on most leucocytes including monocytes, granulocytes, lymphocytes, NK cells, bone marrow myeloid progenitor cells and on a subset of thymocytes.			
	CD62L plays an important role in leucocyte tethering and rolling on the endothelial cell surface and for the homing of naïve lymphocytes to lymph nodes and Peyers patches via HEV. Neutrophils require a constant supply of this molecule on the cell surface for migration into peripheral tissues and adhesion to activated endothelium at sites of inflammation, where CD62L is rapidly shed as soluble L-selectin, but surface expression still remains.			
	The expression of CD62L is down regulated on lymphocytes and neutrophils by PMA stimulation.			
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.			
References	<ol> <li>Zola, H. <i>et al.</i> (1991) The expression of sub-population markers on B cells: a re-evaluation using high-sensitivity fluorescence flow cytometry. <u>Dis Markers. 9 (2):</u> <u>103-18.</u></li> <li>Sopp, P. &amp; Howard, C.J. (1997) Cross-reactivity of monoclonal antibodies to defined human leucocyte differentiation antigens with bovine cells. <u>Vet Immunol Immunopathol. 56</u> (<u>1-2</u>): <u>11-25.</u></li> <li>Haanstra, K.G. <i>et al.</i> (2008) Characterization of naturally occurring CD4+CD25+ regulatory T cells in rhesus monkeys. <u>Transplantation 85:1185-92.</u></li> </ol>			

	<ul> <li>4. Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. <u>Blood. 112 (6): 2512-9.</u></li> <li>5. Raposo, R.A. <i>et al.</i> (2011) Protein Kinase C and NF-{kappa}B-Dependent CD4 Downregulation in Macrophages Induced by T Cell-Derived Soluble Factors:</li> </ul>
	Consequences for HIV-1 Infection. <u>J Immunol. 187: 748-59.</u> 6. Hughes, S.F. <i>et al.</i> (2010) Total hip and knee replacement surgery results in changes in leukocyte and endothelial markers. <u>J Inflamm (Lond). 7:2.</u>
	<ol> <li>7. Bismarck, D. <i>et al.</i> (2012) Canine CD4+CD8+ double positive T cells in peripheral blood have features of activated T cells. <u>Vet Immunol Immunopathol. 149: 157-66.</u></li> <li>8. Hartley, A.N. &amp; Tarleton, R.L. (2015) Chemokine receptor 7 (CCR7)-expression and</li> </ol>
	IFNγ production define vaccine-specific canine T-cell subsets. <u>Vet Immunol Immunopathol.</u> <u>164 (3-4): 127-36.</u>
	9. Hayhoe, R.P. <i>et al.</i> (2006) Annexin 1 and its bioactive peptide inhibit neutrophil- endothelium interactions under flow: indication of distinct receptor involvement. <u>Blood. 107</u> (5): 2123-30.
	<ul> <li>(5). 2123-30.</li> <li>10. Urquhart, P. <i>et al.</i> (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. <u>J Pharmacol Exp Ther. 321 (2): 656-62.</u></li> <li>11. Aspinall, A.I. <i>et al.</i> (2010) CX(3)CR1 and vascular adhesion protein-1-dependent</li> </ul>
	recruitment of CD16(+) monocytes across human liver sinusoidal endothelium. Hepatology. 51 (6): 2030-9.
	12. Rothe, K. <i>et al.</i> (2017) Canine peripheral blood CD4 <sup>+</sup> CD8 <sup>+</sup> double-positive Tcell subpopulations exhibit distinct Tcell phenotypes and effector functions. <u>Vet Immunol</u>
	Immunopathol. 185: 48-56. 13. Withers, S.S. <i>et al.</i> (2018) Multi-color flow cytometry for evaluating age-related changes in memory lymphocyte subsets in dogs. <u>Dev Comp Immunol. 87: 64-74.</u>
	14. Hughes, S.F. <i>et al.</i> (2020) The role of phagocytic leukocytes following flexible ureterenoscopy, for the treatment of kidney stones: an observational, clinical pilots-study. <u>Eur J Med Res. 25 (1): 68.</u>
	15. Svitek, N. <i>et al.</i> (2018) An Ad/MVA vectored <i>Theileria parva</i> antigen induces schizont- specific CD8 <sup>+</sup> central memory T cells and confers partial protection against a lethal challenge. <u>NPJ Vaccines. 3: 35.</u>
	16. Tucker, N. <i>et al.</i> (2023) Bovine blood and milk T-cell subsets in distinct states of activation and differentiation during subclinical <i>Staphylococcus aureus</i> mastitis. <u>J Reprod</u> <u>Immunol. 156: 103826.</u>
	17. Yamauchi, A. <i>et al.</i> (2023) Negative Influence of Aging on Differentiation and Proliferation of CD8(+) T-Cells in Dogs. <u>Vet Sci. 10 (9): 541</u>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts.
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1076SBB675

#### Regulatory

For research purposes only

### **Related Products**

### **Recommended Useful Reagents**

### HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M408042:221010'

#### Printed on 11 Jul 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint