

Datasheet: MCA1079F

Description:	MOUSE ANTI HORSE CD5:FITC
Specificity:	CD5
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	CVS5
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10
Immunofluorescence			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Horse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Equine leucocytes.		
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the mouse X63-Ag8.653 myeloma cell line.		
Specificity	Mouse anti Horse CD5 antibody, clone CVS5 recognizes the equine CD5 antigen, a single-pass		

type I membrane protein with an approximate molecular weight of 69 kDa. Equine CD5 is expressed on the majority of T-lymphocytes, in addition it has been reported that equine CD5 may also be detected at very low levels on B-cells and granulocytes.

In addition to the CVS5 clone, other [CVS clones](#) recognising equine MHC and cell surface antigens are available from Bio-Rad.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

- References**
1. Lunn, D.P. *et al.* (1991) Three monoclonal antibodies identifying antigens on all equine T lymphocytes, and two mutually exclusive T-lymphocyte subsets. [Immunology. 74 \(2\): 251-7.](#)
 2. Lunn, D.P. *et al.* (1998) Report of the Second Equine Leucocyte Antigen Workshop, Squaw valley, California, July 1995. [Vet Immunol Immunopathol. 62 \(2\): 101-43.](#)
 3. Moyo, N.A. *et al.* (2013) Differentiation and activation of equine monocyte-derived dendritic cells are not correlated with CD206 or CD83 expression. [Immunology. 139 \(4\): 472-83.](#)
 4. Mayall, S. *et al.* (2001) The anti-human CD21 antibody, BU33, identifies equine B cells. [J Comp Pathol. 124 :83-7.](#)
 5. Siedek, E. *et al.* (1997) Isolation and characterisation of equine dendritic cells. [Vet Immunol Immunopathol. 60: 15-31.](#)
 6. Colbath AC *et al.* (2016) Autologous and Allogeneic Equine Mesenchymal Stem Cells Exhibit Equivalent Immunomodulatory Properties *In Vitro*. [Stem Cells Dev. Dec 13. \[Epub ahead of print\]](#)
 7. Ziegler, A. *et al.* (2016) Equine dendritic cells generated with horse serum have enhanced functionality in comparison to dendritic cells generated with fetal bovine serum. [BMC Vet Res. 12 \(1\): 254.](#)
 8. Ziegler, A. *et al.* (2016) Identification and characterization of equine blood plasmacytoid dendritic cells. [Dev Comp Immunol. 65: 352-7.](#)
-

Storage Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M364755:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)