

Datasheet: MCA1086F

| | |
|----------------------|---|
| Description: | MOUSE ANTI HORSE MHC CLASS I MONOMORPHIC:FITC |
| Specificity: | MHC CLASS I MONOMORPHIC |
| Format: | FITC |
| Product Type: | Monoclonal Antibody |
| Clone: | CVS22 |
| Isotype: | IgG2a |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | Neat - 1/10 |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | | | |
|--------------------------------|---|---------------------|-------------------|
| Target Species | Horse | | |
| Product Form | Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
| | FITC | 490 | 525 |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant | | |
| Buffer Solution | Phosphate buffered saline | | |
| Preservative | 0.09% sodium azide (NaN ₃) | | |
| Stabilisers | 1% bovine serum albumin | | |
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml | | |
| Immunogen | Equine leucocytes. | | |

| | |
|--------------------------------------|--|
| Fusion Partners | Spleen cells from immunized mice were fused with cells of the X63.Ag 8.653 mouse myeloma cell line. |
| Specificity | <p>Mouse anti Horse MHC Class I Monomorphic antibody, clone CVS22 recognizes monomorphic equine MHC Class I and was classified at the International Equine Leucocyte Antigen Workshop. MHC class I is expressed by all nucleated cells.</p> <p>The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In horses, this complex is referred to as the equine leukocyte antigen (ELA) region.</p> |
| Flow Cytometry | Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl |
| References | <ol style="list-style-type: none"> 1. Lunn, D.P. <i>et al.</i> (1998) Report of the Second Equine Leucocyte Antigen Workshop, Squaw valley, California, July 1995. Vet Immunol Immunopathol. 62:101-143 2. Mérand, C. <i>et al.</i> (2009) Young foal and adult horse monocyte-derived dendritic cells differ by their degree of phenotypic maturity. Vet Immunol Immunopathol. 131 (1-2): 1-8. 3. Carrade, D.D. <i>et al.</i> (2011) Clinicopathologic findings following intra-articular injection of autologous and allogeneic placentally derived equine mesenchymal stem cells in horses. Cytotherapy. 13 (4): 419-30. 4. Soboll Hussey, G. <i>et al.</i> (2014) Innate immune responses of airway epithelial cells to infection with equine herpesvirus-1. Vet Microbiol. 170 (1-2): 28-38. 5. Tessier, L. <i>et al.</i> (2015) Phenotypic and immunomodulatory properties of equine cord blood-derived mesenchymal stromal cells. PLoS One. 10 (4): e0122954. 6. Maumus M <i>et al.</i> (2016) Utility of a Mouse Model of Osteoarthritis to Demonstrate Cartilage Protection by IFNγ-Primed Equine Mesenchymal Stem Cells. Front Immunol. 7: 392. 7. Barberini, D.J. <i>et al.</i> (2018) Safety and tracking of intrathecal allogeneic mesenchymal stem cell transplantation in healthy and diseased horses. Stem Cell Res Ther. 9 (1): 96. 8. Kamm, J.L. <i>et al.</i> (2021) Immune response to allogeneic equine mesenchymal stromal cells. Stem Cell Res Ther. 12 (1): 570. 9. Rapacz-leonard, A. <i>et al.</i> (2018) Major histocompatibility complex class I in the horse (<i>Equus caballus</i>) placenta during pregnancy and parturition. Placenta. 74: 36-46. |
| Storage | <p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p> |
| Guarantee | 12 months from date of despatch |
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1086F 10041 |
| Regulatory | For research purposes only |

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M408078:221010'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)