

Datasheet: MCA1095GA BATCH NUMBER 161500

Description:	ption: MOUSE ANTI SHEEP CD41/CD		
Specificity:	CD41/CD61		
Other names:	GPIIb-IIIa		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	CO.35E4		
lsotype:	lgG1		
Quantity:	0.1 mg		

Product Details

Applications	This product has been reported to work in the following applications. This information is						
	•	derived from testing within our laboratories, peer-reviewed publications or personal					
	communications from the	-					
	information. For general	information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	•			1/50 - 1/200		
	Immunohistology - Frozen						
	Immunohistology - Paraffin						
	ELISA	•					
	Immunoprecipitation	•					
	Western Blotting						
	Where this antibody has necessarily exclude its u a guide only. It is recomm	se in suc	h procedu	res. Suggested workin	g dilutions are given as		
	necessarily exclude its u	se in suc nended tl	h procedu nat the us	res. Suggested workin er titrates the antibody	g dilutions are given as		
Target Species	necessarily exclude its us a guide only. It is recomm	se in suc nended tl	h procedu nat the us	res. Suggested workin er titrates the antibody	g dilutions are given as		
Target Species Species Cross	necessarily exclude its us a guide only. It is recomm system using appropriate	se in suc nended tl e negative	h procedu nat the us e/positive	res. Suggested workin er titrates the antibody controls.	g dilutions are given as		
	necessarily exclude its u a guide only. It is recomm system using appropriate Sheep	se in suc nended th e negative Rabbit, De	h procedu nat the us e/positive	res. Suggested workin er titrates the antibody controls.	g dilutions are given as for use in his/her own		
Species Cross	necessarily exclude its us a guide only. It is recommon system using appropriate Sheep Reacts with: Goat, Cat, F	se in suc nended tl e negative Rabbit, De	h procedu nat the us e/positive og, Horse ng conditi	res. Suggested workin er titrates the antibody controls. , Bovine ons may vary betweer	ng dilutions are given as for use in his/her own		
Species Cross	necessarily exclude its us a guide only. It is recommon system using appropriate Sheep Reacts with: Goat, Cat, F N.B. Antibody reactivity a	se in suc nended th e negative Rabbit, De and worki testing w	h procedu nat the us e/positive og, Horse ng conditi ithin our la	res. Suggested workin er titrates the antibody controls. , Bovine ons may vary betweer aboratories, peer-revie	ng dilutions are given as for use in his/her own n species. Cross wed publications or		
Species Cross	necessarily exclude its us a guide only. It is recommon system using appropriate Sheep Reacts with: Goat, Cat, F N.B. Antibody reactivity a reactivity is derived from	se in suc nended th e negative Rabbit, De and worki testing w	h procedu nat the us e/positive og, Horse ng conditi ithin our la	res. Suggested workin er titrates the antibody controls. , Bovine ons may vary betweer aboratories, peer-revie	ng dilutions are given as for use in his/her own n species. Cross wed publications or		
Species Cross	necessarily exclude its us a guide only. It is recommon system using appropriate Sheep Reacts with: Goat, Cat, F N.B. Antibody reactivity a reactivity is derived from personal communication	se in suc nended th e negative Rabbit, De and worki testing w	h procedu nat the us e/positive og, Horse ng conditi ithin our la	res. Suggested workin er titrates the antibody controls. , Bovine ons may vary betweer aboratories, peer-revie	ng dilutions are given as for use in his/her own n species. Cross wed publications or		

	supernatant	
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)	
Carrier Free	Yes	
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	Sheep leucocytes.	
Fusion Partners	Spleen cells from immunised mice were fused the cells of the myeloma cell line.	SP2-0/Ag14 mouse
Specificity	Mouse anti Sheep CD41/CD61 antibody, clone CO.35E4 real CD41/CD61, also known as GpIIb/IIIa. CD41/CD61 is a calcius heterodimer and is expressed almost exclusively on platelets a megakaryocytes. This complex is an activation-dependent rec Asp)-containing glycoproteins such as fibrinogen, fibronectin, w thrombospondin and vitronectin. It plays a central role in platel haemostasis.	m-complex dependent and on their precursors, the eptor for RGD (Arg-Gly- von Willebrand factor,
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in	n 100ul.
References	 Brodersen, R. <i>et al.</i> (1998) Analysis of the immunological of characterized monoclonal antibodies with specificities against antigens of human and 11 animal species. <u>Vet Immunol Immu</u> Pintado, C.O. <i>et al.</i> (1995) Characterisation of a membrane equine platelets and peripheral blood leukocytes similar to the glycoprotein IIb/IIIa (CD41/61). <u>Vet Immunol Immunopathol. 44</u> Lalko, C.C. <i>et al.</i> (2003) Equine platelet CD62P (P-selectin) and morphologic study. <u>Vet Immunol Immunopathol. 91 (2): 11</u> Major, T.C. <i>et al.</i> (2010) The attenuation of platelet and mor model of extracorporeal circulation by a nitric oxide releasing p<u>2736-45.</u> Brainard, B.M. <i>et al.</i> (2012) Treatment with aspirin or clopid platelet expression of P selectin or platelet-neutrophil aggrega <u>Immunopathol. 149: 119-25.</u> Iwaszko-Simonik, A. <i>et al.</i> (2015) Expression of surface plat CD41/61) in horses with recurrent airway obstruction (RAO). <u>164 (1-2): 87-92.</u> Finding, E.J.T. <i>et al.</i> (2018) Cell specific microvesicles vary predisposition in healthy and previously laminitic ponies. <u>Vet Immunolation in healthy and previously laminitic ponies.</u> <u>Vet Immunolation in healthy and previously laminitic ponies.</u> <u>Vet Immunolation in the altereteres.</u> 	various leucocyte surface nopathol. 64 (1): 1-13. receptor on ruminants and human integrin receptor 4 (3-4): 359-68. expression: a phenotypic 9-34. nocyte activation in a rabbit polymer. <u>Biomaterials. 31:</u> ogrel does not affect equine tes. <u>Vet Immunol</u> telet receptors (CD62P and <u>/et Immunol Immunopathol.</u> with season and disease <u>mmunol Immunopathol. 202:</u>

	equine systemic inflammatory response syndrome. <u>J Vet Diagn</u> <u>10406387221077969.</u>	Invest.	
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.		
	Avoid repeated freezing and thawing as this may denature the a frost-free freezers is not recommended.	intibody. Storage in	
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1095GA 10040		
Regulatory	For research purposes only		

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	RPE
Goat Anti Mouse IgG (STAR70)	FITC
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos., DyLight®488, DyLight®550,</u>
	DyLight®650, DyLight®680, DyLight®800,
	FITC, HRP
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	FITC
Goat Anti Mouse IgG (STAR77)	HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-	rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M395219:220324'

Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint