

Datasheet: MCA1108G

BATCH NUMBER 159859

Description:	escription: RAT ANTI MOUSE CD8	
Specificity:	CD8 ALPHA	
Other names:	LY-2	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	YTS105.18	
Isotype:	IgG2a	
Quantity:	0.25 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/50 - 1/100
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA			•	
Immunoprecipitation				
Western Blotting				
Immunofluorescence	•			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide

Stabilisers Carrier Free Yes Approx. Protein IgG concentration 1 mg/ml Concentrations Immunogen Mouse spleen cells. **External Database** UniProt: Links P01731 Related reagents **Entrez Gene:** 12525 Cd8a Related reagents **Synonyms** Lyt2, Lyt-2 **RRID** AB 322819 **Fusion Partners** Spleen cells from an immunised DA rat were fused with cells of the rat Y3/Ag1.2.3. myeloma cell line. **Specificity** Rat anti Mouse CD8, clone YTS105.18 recognizes a non polymorphic epitope on the mouse CD8 alpha chain. This antibody has been reported to block MHC I dependent T cell responses in vitro and in vivo, and induces transplantation tolerance in combination with CD4 antibodies (Cobbold et al. 1990 & Wise et al. 1998). Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul. References 1. Qin, S.X. et al. (1990) Induction of tolerance in peripheral T cells with monoclonal antibodies. Eur J Immunol. 20 (12): 2737-45. 2. Cobbold, S.P. et al. (1990) The induction of skin graft tolerance in major histocompatibility complex-mismatched or primed recipients; primed T cells can be tolerized in the periphery with anti-CD4 and anti-CD8 antibodies. Eur J Immunol. 20 (12): 2747-55. indirect recognition. J Immunol. 161 (11): 5813-6.

- 3. Wise, M.P. et al. (1998) Linked suppression of skin graft rejection can operate through
- 4. Lacroix-Lamande, S. et al. (2009) Neonate intestinal immune response to CpG oligodeoxynucleotide stimulation. PLoS One. 4: 1-8.
- 5. Auray, G. et al. (2007) Involvement of intestinal epithelial cells in dendritic cell recruitment during C. parvum infection Microbes Infect. 9: 574-82.
- 6. Sroga, J.M. et al. (2003) Rats and mice exhibit distinct inflammatory reactions after spinal cord injury. J Comp Neurol. 462: 223-40.
- 7. Karlsson, M.R. et al. (2010) Hypersensitivity and oral tolerance in the absence of a secretory immune system. Allergy. 65: 561-70.
- 8. Himoudi, N. et al. (2007) Development of anti-PAX3 immune responses; a target for cancer immunotherapy Cancer Immunol Immunother. 56: 1381-95.
- 9. Nakashima, H. et al. (2011) A Novel Combination Immunotherapy for Cancer by

IL-13Rα2-Targeted DNA Vaccine and Immunotoxin in Murine Tumor Models. <u>J Immunol.</u> 187: 4935-46.

- 10. Shaw, T.N. *et al.* (2015) Perivascular Arrest of CD8+ T Cells Is a Signature of Experimental Cerebral Malaria. <u>PLoS Pathog. 11 (11): e1005210.</u>
- 11. Jalili, R.B. *et al.* (2018) Fibroblast cell-based therapy prevents induction of alopecia areata in an experimental model. <u>Cell Transplant. 27 (6): 994-1004.</u>
- 12. Zhao, Q. *et al.* (2019) Tumor-targeted IL-12 combined with tumor resection yields a survival-favorable immune profile. <u>J Immunother Cancer. 7 (1): 154.</u>
- 13. Mohanta, S.K. *et al.* (2022) Neuroimmune cardiovascular interfaces control atherosclerosis. <u>Nature</u>. <u>Apr 27 [Epub ahead of print]</u>.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1108G 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...) DyLight®800

Rabbit Anti Rat IgG (STAR17...)

Goat Anti Rat IgG (STAR72...)

Goat Anti Rat IgG (STAR69...)

Goat Anti Rat IgG (STAR73...)

Rabbit Anti Rat IgG (STAR21...)

HRP

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) DyLight®650, DyLight®650, DyLight®650, DyLight®650, DyLight®650, DyLight®650, DyLight®650, DyLight@650, <a href="mailto:DyLig

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

Recommended Negative Controls

RAT IgG2a NEGATIVE CONTROL (MCA1212)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M382907:210513'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint