Datasheet: MCA1114 BATCH NUMBER 164590

Description:	MOUSE ANTI HUMAN CD45RB
Specificity:	CD45RB
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MEM-55
lsotype:	lgG1
Quantity:	0.2 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry				1/50 - 1/100	
	Immunohistology - Frozen	•				
	Immunohistology - Paraffin	-			1/50 - 1/200	
	ELISA			•		
	Immunoprecipitation	•				
	Western Blotting					
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.					
Target Species	Human					
Species Cross Reactivity	Reacts with: Rhesus Monkey, Monkey Does not react with:Horse, Goat, Llama N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.			ewed publications or		
Product Form	Purified IgG - liquid					
Preparation	Purified IgG prepared by	affinity ch	nromatogi	aphy on Protein A fror	m tissue culture	

	supernatant			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% sodium azide (NaN ₃)			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
External Database Links	UniProt: P08575 Related reagents Entrez Gene: 5788 PTPRC Related reagents			
Synonyms	CD45			
RRID	AB_321430			
Specificity	Mouse anti Human CD45RB antibody, clone MEM-55 recognizes the isoform of the leucocyte common antigen encoded by exon B and stains B cells, most T cells and monocytes. Mouse anti Human CD45RB antibody, clone MEM-55 immunoprecipitates bands of 190, 200 and 205 kDa.			
Flow Cytometry	Use 10µl of the suggested working dilution to label 10^6 cells in $100µ$ l			
References	 Blaylock, M.G. <i>et al.</i> (2003) Eosinophils from patients with asthma express higher level of the pan-leucocyte receptor CD45 and the isoform CD45RO. <u>Clin Exp Allergy. 33</u>: <u>936-41</u>. Windhagen, A. <i>et al.</i> (2007) Altered CD45 isoform expression in C77G carriers influences cytokine responsiveness and adhesion properties of T cells. <u>Clin Exp Immuno</u>. <u>150</u>: 509-17. Ibrahim, S. <i>et al.</i> (2007) Screening of anti-human leukocyte monoclonal antibodies for reactivity with equine leukocytes. <u>Vet Immunol Immunopathol.</u> <u>119</u>: 63-80. Davis, W.C. <i>et al.</i> (2007) Use of flow cytometry to identify monoclonal antibodies that recognize conserved epitopes on orthologous leukocyte differentiation antigens in goats, llamas, and rabbits. <u>Vet Immunol Immunopathol.</u> <u>119</u>: <u>123-30</u>. Ponchel, F. <i>et al.</i> (2014) An immunological biomarker to predict MTX response in early RA. <u>Ann Rheum Dis.</u> <u>73</u> (<u>11</u>): <u>2047-53</u>. Khan, A.A. <i>et al.</i> (2015) Therapeutic immunization with a mixture of herpes simplex virus 1 glycoprotein D-derived "asymptomatic" human CD8+ T-cell epitopes decreases spontaneous ocular shedding in latently infected HLA transgenic rabbits: association witl low frequency of local PD-1+ TIM-3+ CD8+ exhausted T cells. <u>J Virol.</u> <u>89</u> (<u>13</u>): <u>6619-32</u>. Hunt, L. <i>et al.</i> (2016) T cell subsets: an immunological biomarker to predict progression to clinical arthritis in ACPA-positive individuals. <u>Ann Rheum Dis.</u> <u>75</u> (<u>10</u>): <u>1884-9</u>. 	1 <u>1.</u> /		

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1114 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77)	HRP			
Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,			
	DyLight®650, DyLight®680, DyLight®800,			
	FITC, HRP			
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>			
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP			
Recommended Negative Controls				

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M410484:221028'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint