

# Datasheet: MCA1133PE

Description:	MOUSE ANTI HUMAN CD49a:RPE
Specificity:	CD49a
Other names:	INTEGRIN ALPHA 1 CHAIN, VLA-1
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	TS2/7
Isotype:	lgG1
Quantity:	100 TESTS

## **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjuga	ated to R. Phycoerythrin	(RPE) - lyophilized
econstitution	Reconstitute with 1.0	0 ml distilled water	
/lax Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	RPE 488nm laser	496	578
reparation	Purified IgG prepare supernatant	ed by affinity chromatogr	raphy on Protein A
fer Solution	Phosphate buffered	saline	
servative	0.09% sodium azide	e (NaN <sub>3</sub> )	
	0.0070000000000000000000000000000000000	( 0)	
tabilisers	1% bovine serum all	( 0,	

lm	mı	un	OC	ien	ì

Human CTL line directed against HLA-DR.

## External Database Links

#### **UniProt:**

P56199 Related reagents

#### **Entrez Gene:**

3672 ITGA1 Related reagents

#### **RRID**

AB 566795

### **Fusion Partners**

Spleen cells from immunized Balb.c mice were fused with cells of the NS1 mouse myeloma.

#### **Specificity**

Mouse anti Human CD49a antibody, clone TS2/7 recognizes the human alpha 1 integrin sub-unit, which forms the VLA-1 heterodimer in association with the beta 1 integrin. VLA-1 is a receptor for collagen and laminin, and is expressed by chronically activated T cells, melanoma cells and smooth muscle cells.

Mouse anti Human CD49a antibody, clone TS2/7 has been reported as being suitable for use in Western Blotting.

Mouse anti Human CD49a antibody, clone TS2/7 is routinely tested in flow cytometry using HeLa cells.

#### Flow Cytometry

Use 10µl of the suggested working dilution to label 10<sup>6</sup> cells in 100µl.

#### References

- 1. Hemler, M.E. *et al.* (1985) Biochemical characterization of VLA-1 and VLA-2. Cell surface heterodimers on activated T cells. J Biol Chem. 260 (28): 15246-52.
- 2. Hafler, D.A. *et al.* (1985) Investigation of in vivo activated T cells in multiple sclerosis and inflammatory central nervous system diseases. <u>Clin Immunol Immunopathol. 37 (2):</u> 163-71.
- 3. Klemmt, P.A. *et al.* (2007) Endometrial cells from women with endometriosis have increased adhesion and proliferative capacity in response to extracellular matrix components: towards a mechanistic model for endometriosis progression. <u>Hum Reprod.</u> 22: 3139-47.
- 4. Berry, P.A. *et al.* (2011) The calcium sensor STIM1 is regulated by androgens in prostate stromal cells. <u>Prostate</u>. 71: 1646-55.
- 5. Elshaw, S.R. *et al.* (2001) A comparison of ocular melanocyte and uveal melanoma cell invasion and the implication of alpha1beta1, alpha4beta1 and alpha6beta1 integrins. <u>Br J Ophthalmol. 85: 732-8.</u>
- 6. Cervar-Zivkovic, M. *et al.* (2011) Endothelin-1 Stimulates Proliferation of First-Trimester Trophoblasts via the A- and B-Type Receptor and Invasion via the B-Type Receptor. <u>J Clin Endocrinol Metab.</u> 96: 3408-15.
- 7. Kaneko, Y. *et al.* (2012) Integrin  $\alpha 1/\beta 1$  and  $\alpha 2/\beta 1$  as a receptor for IgA1 in human glomerular mesangial cells in IgA nephropathy. <u>Int Immunol. 24: 219-32.</u>
- 8. Majali-Martinez, A. *et al.* (2018) Temporal changes of the endothelin system in human cytotrophoblasts during the first trimester of pregnancy. <u>Physiol Res. 67 (Suppl 1):</u>

S247-S255.

9. Zhuang, L. *et al.* (2021) Dermal Pericytes Exhibit Declined Ability to Promote Human Skin Regeneration with Ageing in 3D Organotypic Culture Models <u>Cells. 10 (11): 3051.</u>

Storage Store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend

microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #20487 available at:

https://www.bio-rad-antibodies.com/SDS/MCA1133PE

20487

**Regulatory** For research purposes only

## Related Products

### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

## **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

Fax: +1 919 878 3751

North & South Tel: +1 800 265 7376

America

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419366:230616'

#### Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint