

# Datasheet: MCA1143A647T

Description:	RAT ANTI MOUSE CD40:Alexa Fluor® 647
Specificity:	CD40
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	3/23
Isotype:	lgG2a
Quantity:	25 TESTS/0.25ml

### **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www bio-						
	rad-antibodies com/protocols						
	Yes No Not Determined Suggested Diluti						
	Flow Cytometry	•			Neat - 1/10		
	Where this antibody ha	as not been te	ested for u	se in a particular	technique this does not		
	necessarily exclude its	use in such	procedure	s. It is recommen	ded that the user titrates		
	the antibody for use in	their own sys	stem using	appropriate nega	ative/positive controls.		
Target Species	Mouse						
Product Form	Purified IgG conjugate	d to Alexa Flu	ıor® 647 -	liquid			
Max Ex/Em	Fluorophore	Excitation Ma	ax (nm) E	mission Max (nm)	)		
	Alexa Fluor®647	650		665			
Preparation	Purified IgG prepared supernatant	by affinity chr	omatograp	ohy on Protein G t	from tissue culture		
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum A	Albumin					
Approx. Protein Concentrations	IgG concentration 0.05	i mg/ml					
Immunogen	Extracellular Domain c	of Mouse CD4	0 and the	Fc portion of Hur	nan IgG1.		

External Database Links	UniProt:         P27512       Related reagents         Entrez Gene:         21939       Cd40       Related reagents
Synonyms	Tnfrsf5
RRID	AB_1102005
Fusion Partners	Spleen cells from immunised LOU/c rats were fused with cells of the Ag8 mouse myeloma cell line.
Specificity	<b>Rat anti Mouse CD40 antibody, clone 3/23</b> recognizes the murine CD40 cell surface glycoprotein. It does not react with normal mouse Ig or with human IgG1 and will stain most mature mouse B cells. It does not cross react with mouse T cells. The specificity of Rat anti Mouse CD40 antibody, clone 3/23 was demonstrated by ELISA and flow cytometry using BHK cells transfected with mouse CD40.
	Rat anti Mouse CD40 antibody, clone 3/23 is a powerful activator of normal B cells especially in the presence of IL-4.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ( <u>BUF041A/B</u> ).
References	<ol> <li>Parry, S.L. <i>et al.</i> (1994) Plastic-immobilized anti-mu or anti-delta antibodies induce apoptosis in mature murine B lymphocytes. <u>Eur J Immunol. 24 (4): 974-9.</u></li> <li>Hasbold, J. <i>et al.</i> (1994) Properties of mouse CD40: cellular distribution of CD40 and B cell activation by monoclonal anti-mouse CD40 antibodies. <u>Eur J Immunol. 24 (8):</u> 1835-42.</li> <li>Hasbold, J. &amp; Klaus, G.G. (1994) B cells from CBA/N mice do not proliferate following ligation of CD40. <u>Eur J Immunol. 24 (1): 152-7.</u></li> <li>Bedoret, D. <i>et al.</i> (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. <u>J Clin Invest. 119 (12): 3723-38.</u></li> <li>Mohan, J. <i>et al.</i> (2005) Skin-derived dendritic cells acquire and degrade the scrapie agent following in vitro exposure. <u>Immunology. 116: 122-33.</u></li> <li>Mukhopadhyay, S. <i>et al.</i> (2004) Activation of murine macrophages by Neisseria meningitidis and IFN-gamma in vitro: distinct roles of class A scavenger and Toll-like pattern recognition receptors in selective modulation of surface phenotype. <u>J Leukoc Biol.</u> 76: 577-84.</li> <li>Russo, S. <i>et al.</i> (2003) Platelet-activating factor mediates CD40-dependent angiogenesis and endothelial-smooth muscle cell interaction. <u>J Immunol. 171: 5489-97.</u> 8. Shakib, S. <i>et al.</i> (2009) Checkpoints in the development of thymic cortical epithelial cells. <u>J Immunol. 182: 130-7.</u></li> <li>Ogasawara, K. <i>et al.</i> (2002) Profiles of cell-to-cell interaction of Mycobacterium</li> </ol>

Regulatory	For research purposes only
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</u>
Acknowledgements	This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com
Guarantee	18 months from date of despatch.
	Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
	Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.
	This product should be stored undiluted.
Storage	Store at +4°C or at -20°C if preferred.
	14. Malada-Edelstein, Y.F. <i>et al.</i> (2017) Regulatory role of cytosolic phospholipase A <sub>2</sub> alpha in the induction of CD40 in microglia. <u>J Neuroinflammation. 14 (1): 33.</u>
	cells into Foxp3- IL-10+ regulatory T cells by engaging CD28 and CTLA-4. Eur J Immunol.
	Cells. <u>J Immunol. 187: 4517-29.</u> 13. Pletinckx, K. <i>et al.</i> (2015) Immature dendritic cells convert anergic nonregulatory T
	12. Legutko, A. <i>et al.</i> (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor-{gamma} Activity in Dendritic
	collaboration between v(alpha)14(+) NKT cells and antigen-presenting cells. Int Immunol. <u>12: 1669-75.</u>
	11. Yang, Y.F. <i>et al.</i> (2000) Requirement for IFN-gamma in IL-12 production induced by
	different origins and dynamics. <u>J Immunol. 160: 708-17.</u>
	suppressor signal transmission. <u>Clin Exp Immunol. 129: 272-80.</u> 10. Salomon, B. <i>et al.</i> (1998) Three populations of mouse lymph node dendritic cells with
	intracellulare-induced immunosuppressive macrophages with target T cells in terms of

## **Related Products**

## **Recommended Negative Controls**

RAT IgG2a NEGATIVE CONTROL:Alexa Fluor® 647 (MCA1212A647)

### **Recommended Useful Reagents**

#### MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-rac	d.com	Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M361757:200311'

#### Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint