

## Datasheet: MCA1143GA

<b>Description:</b>	RAT ANTI MOUSE CD40
<b>Specificity:</b>	CD40
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	3/23
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% sodium azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes

<b>Approx. Protein Concentrations</b>	IgG concentration 1 mg/ml
<b>Immunogen</b>	Extracellular Domain of Mouse CD40 and the Fc portion of Human IgG1.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P27512</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">21939</a> Cd40    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Tnfrsf5
<b>RRID</b>	AB_323810
<b>Fusion Partners</b>	Spleen cells from immunised LOU/c rats were fused with cells of the Ag8 mouse myeloma cell line.
<b>Specificity</b>	<p><b>Rat anti Mouse CD40 antibody, clone 3/23</b> recognizes the murine CD40 cell surface glycoprotein. It does not react with normal mouse Ig or with human IgG1 and will stain most mature mouse B cells. It does not cross react with mouse T cells. The specificity of Rat anti Mouse CD40 antibody, clone 3/23 was demonstrated by ELISA and flow cytometry using BHK cells transfected with mouse CD40.</p> <p>Rat anti Mouse CD40 antibody, clone 3/23 is a powerful activator of normal B cells especially in the presence of IL-4.</p>
<b>Flow Cytometry</b>	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Parry, S.L. <i>et al.</i> (1994) Plastic-immobilized anti-mu or anti-delta antibodies induce apoptosis in mature murine B lymphocytes. <a href="#">Eur J Immunol. 24 (4): 974-9.</a></li> <li>2. Hasbold, J. &amp; Klaus, G.G. (1994) B cells from CBA/N mice do not proliferate following ligation of CD40. <a href="#">Eur J Immunol. 24 (1): 152-7.</a></li> <li>3. Salomon, B. <i>et al.</i> (1998) Three populations of mouse lymph node dendritic cells with different origins and dynamics. <a href="#">J Immunol. 160: 708-17.</a></li> <li>4. Yang, Y.F. <i>et al.</i> (2000) Requirement for IFN-gamma in IL-12 production induced by collaboration between v(alpha)14(+) NKT cells and antigen-presenting cells. <a href="#">Int Immunol. 12: 1669-75.</a></li> <li>5. Ogasawara, K. <i>et al.</i> (2002) Profiles of cell-to-cell interaction of Mycobacterium intracellulare-induced immunosuppressive macrophages with target T cells in terms of suppressor signal transmission. <a href="#">Clin Exp Immunol. 129: 272-80.</a></li> <li>6. Russo, S. <i>et al.</i> (2003) Platelet-activating factor mediates CD40-dependent angiogenesis and endothelial-smooth muscle cell interaction. <a href="#">J Immunol. 171: 5489-97.</a></li> <li>7. Mukhopadhyay, S. <i>et al.</i> (2004) Activation of murine macrophages by Neisseria meningitidis and IFN-gamma in vitro: distinct roles of class A scavenger and Toll-like pattern recognition receptors in selective modulation of surface phenotype. <a href="#">J Leukoc Biol. 76: 577-84.</a></li> </ol>

8. Mohan, J. *et al.* (2005) Skin-derived dendritic cells acquire and degrade the scrapie agent following in vitro exposure. [Immunology. 116: 122-33.](#)
9. Bedoret, D. *et al.* (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. [J Clin Invest. 119 \(12\): 3723-38.](#)
10. Shakib, S. *et al.* (2009) Checkpoints in the development of thymic cortical epithelial cells. [J Immunol. 182: 130-7.](#)
11. Legutko, A. *et al.* (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor- $\gamma$  Activity in Dendritic Cells. [J Immunol. 187: 4517-29.](#)
12. Pletinckx, K. *et al.* (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3- IL-10+ regulatory T cells by engaging CD28 and CTLA-4. [Eur J Immunol. 45 \(2\): 480-91.](#)
13. Malada-Edelstein, Y.F. *et al.* (2017) Regulatory role of cytosolic phospholipase A<sub>2</sub> alpha in the induction of CD40 in microglia. [J Neuroinflammation. 14 \(1\): 33.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1143GA>  
10040

---

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight@550</a> , <a href="#">DyLight@650</a> , <a href="#">DyLight@800</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>
Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight@800</a>
Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>
Rabbit Anti Rat IgG (STAR17...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M408304:221011'

**Printed on 04 Sep 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)