

Datasheet: MCA1143PE

Description:	RAT ANTI MOUSE CD40:RPE
Specificity:	CD40
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	3/23
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
	5% Sucrose		
Immunogen	Extracellular Domain of Mouse CD40 and the Fc portion of Human IgG1.		
External Database Links	UniProt: P27512 Related reagents		
	Entrez Gene:		

Synonyms	Tnfrsf5
-----------------	---------

Fusion Partners	Spleen cells from immunised LOU/c rats were fused with cells of the Ag8 mouse myeloma cell line.
------------------------	--

Specificity	<p>Rat anti Mouse CD40 antibody, clone 3/23 recognizes the murine CD40 cell surface glycoprotein. It does not react with normal mouse Ig or with human IgG1 and will stain most mature mouse B cells. It does not cross react with mouse T cells. The specificity of Rat anti Mouse CD40 antibody, clone 3/23 was demonstrated by ELISA and flow cytometry using BHK cells transfected with mouse CD40.</p> <p>Rat anti Mouse CD40 antibody, clone 3/23 is a powerful activator of normal B cells especially in the presence of IL-4.</p>
--------------------	--

Flow Cytometry	<p>Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
-----------------------	--

References	<ol style="list-style-type: none">1. Parry, S.L. <i>et al.</i> (1994) Plastic-immobilized anti-mu or anti-delta antibodies induce apoptosis in mature murine B lymphocytes. Eur J Immunol. 24 (4): 974-9.2. Hasbold, J. <i>et al.</i> (1994) Properties of mouse CD40: cellular distribution of CD40 and B cell activation by monoclonal anti-mouse CD40 antibodies. Eur J Immunol. 24 (8): 1835-42.3. Hasbold, J. & Klaus, G.G. (1994) B cells from CBA/N mice do not proliferate following ligation of CD40. Eur J Immunol. 24 (1): 152-7.4. Bedoret, D. <i>et al.</i> (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. J Clin Invest. 119 (12): 3723-38.5. Mohan, J. <i>et al.</i> (2005) Skin-derived dendritic cells acquire and degrade the scrapie agent following in vitro exposure. Immunology. 116: 122-33.6. Mukhopadhyay, S. <i>et al.</i> (2004) Activation of murine macrophages by Neisseria meningitidis and IFN-gamma in vitro: distinct roles of class A scavenger and Toll-like pattern recognition receptors in selective modulation of surface phenotype. J Leukoc Biol. 76: 577-84.7. Russo, S. <i>et al.</i> (2003) Platelet-activating factor mediates CD40-dependent angiogenesis and endothelial-smooth muscle cell interaction. J Immunol. 171: 5489-97.8. Shakib, S. <i>et al.</i> (2009) Checkpoints in the development of thymic cortical epithelial cells. J Immunol. 182: 130-7.9. Ogasawara, K. <i>et al.</i> (2002) Profiles of cell-to-cell interaction of Mycobacterium intracellulare-induced immunosuppressive macrophages with target T cells in terms of suppressor signal transmission. Clin Exp Immunol. 129: 272-80.10. Salomon, B. <i>et al.</i> (1998) Three populations of mouse lymph node dendritic cells with different origins and dynamics. J Immunol. 160: 708-17.11. Yang, Y.F. <i>et al.</i> (2000) Requirement for IFN-gamma in IL-12 production induced by collaboration between v(alpha)14(+) NKT cells and antigen-presenting cells. Int Immunol. 12: 1669-75.12. Legutko, A. <i>et al.</i> (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor-γ Activity in Dendritic Cells. J Immunol. 187: 4517-29.13. Pletinckx, K. <i>et al.</i> (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3- IL-10+ regulatory T cells by engaging CD28 and CTLA-4. Eur J Immunol. 45 (2): 480-91.14. Malada-Edelstein, Y.F. <i>et al.</i> (2017) Regulatory role of cytosolic phospholipase A₂ alpha in the induction of CD40 in microglia. J Neuroinflammation. 14 (1): 33.
-------------------	---

Storage Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of reconstitution.

Health And Safety Information Material Safety Datasheet documentation #10075 available at:
10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:RPE \(MCA1212PE\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M300867:170106'

Printed on 18 Jul 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)