

Datasheet: MCA1148

BATCH NUMBER 174493

Description:	MOUSE ANTI HUMAN CD71
Specificity:	CD71
Other names:	TRANSFERRIN RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	DF1513
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/25
Immunohistology - Frozen		■		
Immunohistology - Paraffin		■		
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	
Immunofluorescence	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Rhesus Monkey, Mustelid, Ferret N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	KGI cell line.
External Database Links	UniProt: P02786 Related reagents Entrez Gene: 7037 TFRC Related reagents
RRID	AB_1125283
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NSI myeloma cell line.
Specificity	Mouse anti Human CD71 antibody, clone DF1513 recognizes the human CD71 cell surface antigen, a ~190 kDa homodimeric glycoprotein expressed by proliferating cells. CD71 is also known as the transferrin receptor. Mutation of the TFRC gene has been implicated in the development of Immunodeficiency 46 (IMD46) a combined immunodeficiency characterized by early onset chronic diarrhea, recurrent infections and intermittent neutropenia and thrombocytopenia (Jabara <i>et al.</i> 2016).
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl.
References	- Sopper, S. <i>et al.</i> (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. Cytometry. 29 (4): 351-62. - Janes, P.W. <i>et al.</i> (1999) Aggregation of lipid rafts accompanies signaling via the T cell antigen receptor. J Cell Biol. 147: 447-61. - Weissgerber, P. <i>et al.</i> (2003) Investigation of mechanisms involved in phagocytosis of Legionella pneumophila by human cells. FEMS Microbiol Lett. 219 (2): 173-9. - Stockwin, L.H. <i>et al.</i> (2009) Artemisinin dimer anticancer activity correlates with heme-catalyzed reactive oxygen species generation and endoplasmic reticulum stress induction. Int J Cancer. 125: 1266-75. - Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. Vet Immunol Immunopathol. 132:109-15. - Meng, J. <i>et al.</i> (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. PLoS One. 6(3):e17454.

7. Procaccini, C. *et al.* (2012) Leptin-induced mTOR activation defines a specific molecular and transcriptional signature controlling CD4⁺ effector T cell responses. [J Immunol. 189: 2941-53.](#)
8. Makoveichuk, E. *et al.* (2012) Inactivation of lipoprotein lipase occurs on the surface of THP-1 macrophages where oligomers of angiopoietin-like protein 4 are formed. [Biochem Biophys Res Commun. 425:138-43.](#)
9. Trakarnsanga, K. *et al.* (2017) An immortalized adult human erythroid line facilitates sustainable and scalable generation of functional red cells. [Nat Commun. 8: 14750.](#)
10. Deng, K. *et al.* (2023) Application of *In vitro* transcytosis models to brain targeted biologics. [PLoS One. 18 \(8\): e0289970.](#)
11. Grain, A. *et al.* (2025) Evaluation of multi-antigen targeting ADCC strategies in pediatric BCP-ALL. [J Immunother Cancer. 13 \(12\): e012552.](#)

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1148
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M408328:221011'

Printed on 24 Aug 2026

© 2026 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)