

Datasheet: MCA116F

Description:	MOUSE ANTI HUMAN HLA B27:FITC		
Specificity:	HLA B27		
Format:	FITC		
Product Type:	Monoclonal Antibody		
Clone:	HLA-ABC-m3		
Isotype:	lgG2a		
Quantity:	100 TESTS/1ml		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat
Immunohistology - Frozen			•	
Immunohistology - Paraffin				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Human			
Purified IgG conju	gated to Fluorescein Isotl	niocyanate Isomer 1	(FITC) - liquid
Fluorophore	Excitation Max (nm)	Emission Max (nm)	
FITC	490	525	
Purified IgG prepa	ared by affinity chromatog	raphy on Protein A	
Phosphate buffere	ed saline		
0.1% Sodium Azio	de (NaN ₃)		
0.2% Bovine Seru	ım Albumin		
50 ug/ml			
	Purified IgG conjunctions Fluorophore FITC Purified IgG preparation of the properties of the propert	Purified IgG conjugated to Fluorescein Isotle Fluorophore Excitation Max (nm) FITC 490 Purified IgG prepared by affinity chromatog Phosphate buffered saline 0.1% Sodium Azide (NaN ₃) 0.2% Bovine Serum Albumin	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 Fluorophore Excitation Max (nm) Emission Max (nm) FITC 490 525 Purified IgG prepared by affinity chromatography on Protein A Phosphate buffered saline 0.1% Sodium Azide (NaN ₃) 0.2% Bovine Serum Albumin

Immunogen

Immune complex precipitated from an HLA-B27 positive cell line by anti HLA antibody and staphylococcal protein A.

External Database

Links

UniProt:

P03989 Related reagents

Entrez Gene:

3106 HLA-B Related reagents

Synonyms

HLAB

RRID

AB 322098

Specificity

Mouse anti Human HLA B27 antibody, clone HLA-ABC-m3 recognizes the HLA-B27 alloantigen. This antibody reacts with the peripheral blood lymphocytes of 47/47 individuals conventionally typed as HLA-B27 + and precipitates cell surface molecules of 43 and 12 kD, corresponding to the HLA Class 1 heavy chain and beta 2 microglobulin.

Affinity studies by Scatchard analysis showed that this antibody has a higher affinity for HLA-B27 ($9.7 \times 10^8 M^{-1}$) than for HLA B7 ($9.5 \times 10^7 M^{-1}$).

Subjects Expected Fluorescence Intensity Heterozygous HLA-B27+ Strong Homozygous HLA-B27+ Strong Heterozygous HLA-B7+ Faint Non B27, Non B7 Negative

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells or 100ul of whole blood.

References

- 1. Trapani, J.A. *et al.* (1983) Description of a mouse monoclonal anti-HLA-B27 antibody HLA-ABC-m3. <u>Hum Immunol.</u> 7 (4): 205-16.
- 2. Orr, K. *et al.* (1994) Utilization of commercial antisera and flow cytometry in HLA-B27 typing. Cytometry (Comm. Clin. Cytometry) 18: 17-20.
- 3. Levering, W.H. *et al.* (2003) Flow cytometric HLA-B27 screening: cross-reactivity patterns of commercially available anti-HLA-B27 monoclonal antibodies with other HLA-B antigens. Cytometry B Clin Cytom. 54: 28-38.
- 4. Mandic, R. *et al.* (2004) Comparison of surface HLA class I levels in squamous cell carcinoma cell lines of the head and neck. <u>Anticancer Res. 24 (2B): 973-9.</u>
- 5. Goodall, J.C. *et al.* (2007) Does HLA-B27 influence the monocyte inflammatory response to lipopolysaccharide? Rheumatology (Oxford). 46: 232-7.
- 6. Meyer, O. *et al.* (2006) A simple and practical agglutination assay for human leucocyte antigen-B27 typing. Vox Sang. 91 (1): 77-80.
- 7. Meyer, O. *et al.* (2008) Application of the particle gel agglutination assay in the typing of single human leucocyte antigens. <u>Tissue Antigens</u>. 71 (2): 157-9.
- 8. Marroquin, B. O *et al.* (2015) HLA-B27-Homodimer-Specific Antibody Modulates the Expansion of Pro-Inflammatory T-Cells in HLA-B27 Transgenic Rats. <u>PLoS One. 10 (6)</u>: e0130811.

Storage

Store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M364855:200529'

Printed on 22 Mar 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint