

# Datasheet: MCA1187T

MOUSE ANTI HUMAN CD49e
CD49e
INTEGRIN ALPHA 5 CHAIN, VLA-5
Purified
Monoclonal Antibody
JBS5
lgG1

## **Product Details**

					s information is derived		
	from testing within our labor	-					
	•	the originators. Please refer to references indicated for further information. For general protocol					
	recommendations, please v						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	•			Neat - 1/10		
	Immunohistology - Frozen	•			0.5ug/ml - 1ug/ml		
	Immunohistology - Paraffin						
	ELISA						
	Immunoprecipitation	•			1ug/ml - 5ug/ml		
	Western Blotting						
	Where this antibody has no	t been tes	ted for use	in a particular technique	this does not necessarily		
	exclude its use in such proc	edures. S	uggested v	vorking dilutions are give	en as a guide only. It is		
	recommended that the user	<sup>-</sup> titrates th	e antibody	for use in their own syst	em using appropriate		
	negative/positive controls.						
Target Species	Human						
Species Cross	Reacts with: Dog						
Reactivity	<b>N.B.</b> Antibody reactivity and working conditions may vary between species.						
Product Form	Purified IgG - liquid						
Preparation	Purified IgG prepared by affinity chromatography on Protein A.						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide						
Approx. Protein Concentrations	lg concentration 1.0 mg/ml						

External Database Links	UniProt: <u>P08648</u> <u>Related reagents</u> Entrez Gene: <u>3678</u> ITGA5 <u>Related reagents</u>				
Synonyms	FNRA				
RRID	AB_2128198				
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.				
Specificity	<b>Mouse anti Human CD49e antibody, clone JBS5</b> recognizes the Alpha 5 integrin, also known as CD49e, VLA-5, Fibronectin receptor subunit alpha or Integrin alpha-F. CD49e is a 1049 amino acid ~130kDa single pass type-1 transmembrane glycoprotein containing seven <u>FG-GAP</u> repeats. Mouse anti human CD49e antibody, clone JBS5 been shown to stimulate homotypic cellular aggregation of alpha 5 beta 1 positive lymphoid cell lines ( <u>Shen <i>et al.</i> 1991</u> ). It also inhibits the adherence of cells to fibronectin.				
	Removal of sodium azide is recommended prior to use in functional assays.				
Flow Cytometry	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul of whole blood.				
Histology Positive Control Tissue	Human Endothelium				
References	<ol> <li>Shen, C.X. <i>et al.</i> (1991) Antibodies to different members of the beta 1 (CD29) integrins induce homotypic and heterotypic cellular aggregation. <u>Cell Immunol. 138 (1): 216-28.</u></li> <li>McCarty, O.J. <i>et al.</i> (2004) Evaluation of the role of platelet integrins in fibronectin-dependent spreading and adhesion. <u>J Thromb Haemost. 2 (10): 1823-33.</u></li> <li>Bax, D.V. <i>et al.</i> (2004) Integrin alpha5beta1 and ADAM-17 interact <i>in vitro</i> and co-localize in migrating HeLa cells. <u>J Biol Chem. 279 (21): 22377-86.</u></li> <li>Gassmann, P. <i>et al.</i> (2009) CXCR4 regulates the early extravasation of metastatic tumor cells <i>in vivo</i>. <u>Neoplasia. 11: 651-61.</u></li> <li>Yao, H. <i>et al.</i> (2010) Increased potency of the PHSCN dendrimer as an inhibitorof human prostate cancer cell invasion, extravasation, and lung colony formation. <u>Clin Exp Metastasis 27:173-84.</u></li> <li>Gassmann, P. <i>et al.</i> (2010) In vivo tumor cell adhesion in the pulmonary microvasculature is exclusively mediated by tumor cellendothelial cell interaction. <u>BMC Cancer. 10: 177.</u></li> <li>Orazizadeh, M. <i>et al.</i> (2008) CD47 associates with alpha 5 integrin and regulates responses of human articular chondrocytes to mechanical stimulation in an in vitro model. <u>Arthritis Res Ther. 10: R4.</u></li> </ol>				
Storage	Store at +4°C or at -20°C if preferred.				
	This product should be stored undiluted.				
	Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.				

Guarantee	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</u>
Regulatory	For research purposes only

## **Related Products**

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>				
Goat Anti Mouse IgG (STAR77)	HRP			
Rabbit Anti Mouse IgG (STAR12)	RPE			
Rabbit Anti Mouse IgG (STAR8)	DyLight®800			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR76)	RPE			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG (Fc) (STAR120)	<u>FITC</u> , <u>HRP</u>			
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>			
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®680</u> , <u>DyLight®800</u> , <u>FITC</u> , <u>HRP</u>			

### **Recommended Negative Controls**

### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bic	-rad.com	Email: antibody_sales_de@bio-rad.com

'M359757:191004'

#### Printed on 28 Oct 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint