

Datasheet: MCA1193EL

Description:	MOUSE ANTI HUMAN CD16:Low Endotoxin
Specificity:	CD16
Other names:	FcRIII
Format:	Low Endotoxin
Product Type:	Monoclonal Antibody
Clone:	LNK16
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen		▪		
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Functional Assays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by ion exchange chromatography from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Endotoxin Level	<0.01 EU/ug

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Normal human peripheral blood granulocytes.
External Database Links	<p>UniProt:</p> <p>P08637 Related reagents</p> <p>O75015 Related reagents</p> <p>Entrez Gene:</p> <p>2214 FCGR3A Related reagents</p> <p>2215 FCGR3B Related reagents</p>
Synonyms	CD16A, CD16B, FCG3, FCGR3, IGFR3
RRID	AB_566548
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.653 myeloma cell line.
Specificity	<p>Mouse anti Human CD16 antibody, clone LNK16 recognizes CD16, a 50-65 kDa cell surface molecule which exists in two forms - a transmembranous form expressed by NK cells and some T cells, and a phosphatidylinositol linked form expressed by granulocytes.</p> <p>CD16 is a low affinity receptor for IgG (FcR III), and is an important receptor mediating ADCC by NK cells. Mouse anti Human CD16 antibody, clone LNK16 has been reported to block the binding of IgG to the CD16 molecule (Tamm and Schmidt 1996).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells or 100µl human whole blood.
References	<ol style="list-style-type: none"> 1. Tamm, A. & Schmidt, R.E. (1996) The binding epitopes of human CD16 (Fc gamma RIII) monoclonal antibodies. Implications for ligand binding. J Immunol. 157 (4): 1576-81. 2. Lee, S.F. <i>et al.</i> (1999) Cytokine receptor common beta chain as a potential activator of cytokine withdrawal-induced apoptosis. Mol Cell Biol. 19: 7399-409. 3. Hanna, J. <i>et al.</i> (2003) CXCL12 expression by invasive trophoblasts induces the specific migration of CD16- human natural killer cells. Blood. 102:1569-77. 4. Sironi, M. <i>et al.</i> (2006) Differential regulation of chemokine production by Fc gamma receptor engagement in human monocytes: association of CCL1 with a distinct form of M2 monocyte activation (M2b, Type 2). J Leukoc Biol. 80: 342-9. 5. Bowles, J.A. <i>et al.</i> (2006) Anti-CD20 monoclonal antibody with enhanced affinity for CD16 activates NK cells at lower concentrations and more effectively than rituximab. Blood. 108: 2648-54. 6. Kahn, F. <i>et al.</i> (2008) Antibodies against a surface protein of <i>Streptococcus pyogenes</i> promote a pathological inflammatory response. PLoS Pathog. 4: e1000149. 7. Meknache, N. <i>et al.</i> (2009) Human basophils express the glycosylphosphatidylinositol-anchored low-affinity IgG receptor Fc gamma RIIIB (CD16B). J Immunol. 182: 2542-50. 8. Welters, I.D. <i>et al.</i> (2010) Ketamine inhibits transcription factors activator protein 1 and

nuclear factor-kappaB, interleukin-8 production, as well as CD11b and CD16 expression: studies in human leukocytes and leukocytic cell lines. [Anesth Analg. 110: 934-41.](#)

9. Armour, K.L. *et al.* (2010) Expression of human FcγRIIIa as a GPI-linked molecule on CHO cells to enable measurement of human IgG binding. [J Immunol Methods. 354: 20-33.](#)

10. Moalli, F. *et al.* (2010) Role of complement and Fcγ receptors in the protective activity of the long pentraxin PTX3 against *Aspergillus fumigatus*. [Blood. 116: 5170-80.](#)

11. Morris, D.L. *et al.* (2010) Evidence for both copy number and allelic (NA1/NA2) risk at the FCGR3B locus in systemic lupus erythematosus. [Eur J Hum Genet. 18 \(9\): 1027-31.](#)

12. Kamp, V.M. *et al.* (2012) Human suppressive neutrophils CD16bright/CD62Ldim exhibit decreased adhesion. [J Leukoc Biol. 92 \(5\): 1011-20.](#)

13. Weissmüller, S. *et al.* (2012) ICOS-LICOS interaction is critically involved in TGN1412-mediated T-cell activation. [Blood. 119: 6268-77.](#)

14. Pillay, J. *et al.* (2012) A subset of neutrophils in human systemic inflammation inhibits T cell responses through Mac-1. [J Clin Invest. 122: 327-36.](#)

15. Onodera, R. *et al.* (2017) Anti-human neutrophil antigen-1a, -1b, and -2 antibodies in neonates and children with immune neutropenias analyzed by extracted granulocyte antigen immunofluorescence assay. [Transfusion. 57 \(11\): 2586-94.](#)

Further Reading

1. Ravetch, J.V. & Kinet, J.P. (1991) Fc receptors. [Annu Rev Immunol. 9: 457-92.](#)

2. van de Winkel, J.G. & Capel, P.J. (1993) Human IgG Fc receptor heterogeneity: molecular aspects and clinical implications. [Immunol Today. 14 \(5\): 215-21.](#)

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10162 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1193EL>
10162

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG (STAR70...) [FITC](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#), [DyLight®650](#), [DyLight®680](#), [DyLight®800](#),

[FITC](#), [HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin \(MCA928EL\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M408390:221012'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)