

# Datasheet: MCA1194SBUV605

| Description:  | MOUSE ANTI HUMAN CD2:StarBright UltraViolet 605 |  |  |
|---------------|---|--|--|
| Specificity:  | CD2   |  |  |
| Other names:  | LFA-2   |  |  |
| Format:       | StarBright UltraViolet 605                      |  |  |
| Product Type: | Monoclonal Antibody                             |  |  |
| Clone:        | LT2   |  |  |
| Isotype:      | lgG2b   |  |  |
| Quantity:     | 100 TESTS/0.5ml                                 |  |  |
|               |   |  |  |

## **Product Details**

| Applications                | ations This product has been reported to work in the following applications. This inform derived from testing within our laboratories, peer-reviewed publications or perso communications from the originators. Please refer to references indicated for fur information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . |                |      |                   |                    |  |
|-----------------------------|--|----------------|------|-------------------|--------------------|--|
|                             |  | Yes            | No   | Not Determined    | Suggested Dilution |  |
|                             | Flow Cytometry   | •              |      |                   | Neat               |  |
|                             | Where this product has not been tested for use in a particular technique this does not<br>necessarily exclude its use in such procedures. Suggested working dilutions are given as<br>a guide only. It is recommended that the user titrates the product for use in their own<br>system using appropriate negative/positive controls.                                      |                |      |                   |                    |  |
| Target Species              | Human  |                |      |                   |                    |  |
| Species Cross<br>Reactivity | Reacts with: Rhesus Monkey, Chimpanzee, Cynomolgus monkey<br><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross<br>reactivity is derived from testing within our laboratories, peer-reviewed publications or<br>personal communications from the originators. Please refer to references indicated for<br>further information.         |                |      |                   |                    |  |
| Product Form                | Purified IgG conjugated to StarBright UltraViolet 605 - liquid   |                |      |                   |                    |  |
| Max Ex/Em                   | Fluorophore  | Excitation Max | (nm) | Emission Max (nm) |                    |  |
|                             | StarBright UltraViolet<br>605  | 340            |      | 609               |                    |  |
| Preparation                 | Purified IgG prepared by affinity chromatography on Protein A from ascites   |                |      |                   |                    |  |
| Buffer Solution             | Phosphate buffered sa  | aline          |      |                   |                    |  |

| Preservative<br>Stabilisers | 0.09% sodium azide (NaN <sub>3</sub> )<br>1% bovine serum albumin<br>0.1% Pluronic F68<br>0.1% PEG 3350<br>0.05% Tween 20  |                   |
|-----------------------------|--|-------------------|
| Immunogen                   | Normal human peripheral blood lymphocytes.   |                   |
| External Database<br>Links  | UniProt:         P06729       Related reagents         Entrez Gene:         914       CD2       Related reagents   |                   |
| Synonyms                    | SRBC   |                   |
| Fusion Partners             | Spleen cells from immunized BALB/c mice were fused with cells of the mouse X63 myeloma cell line.  | 653               |
| Specificity                 | <ul> <li>Mouse anti Human CD2 antibody, clone LT2 recognizes human CD2, a single classical transmembrane glycoprotein of 50 kDa, expressed by T cells, NK cells and thymood</li> <li>CD2 is a receptor for CD58 (LFA-3), and is important in mediating the interaction be T cells and target cells or antigen presenting cells.</li> <li>Mouse anti Human CD2 antibody, clone LT2 inhibits NK cell mediated cytotoxicity, ConA and PHA mediated blast transformation.</li> </ul>   | cytes.<br>between |
| Flow Cytometry              | Use 5µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.  | 3                 |
| References                  | <ol> <li>Lin, C.W. <i>et al.</i> (2005) CD94 1A transcripts characterize lymphoblastic<br/>lymphoma/leukemia of immature natural killer cell origin with distinct clinical feature<br/><u>Blood. 106 (10): 3567-74.</u></li> <li>Kap, Y.S. <i>et al.</i> (2009) A monoclonal antibody selection for immunohistochemical<br/>examination of lymphoid tissues from non-human primates. <u>J Histochem Cytochem</u><br/><u>1159-67.</u></li> <li>Bruger, A.M. <i>et al.</i> (2020) Protocol to assess the suppression of T-cell proliferation<br/>human MDSC. <u>Methods Enzymol. 632: 155-92.</u></li> </ol> | al<br>n. 57:      |
| Further Reading             | 1. Moingeon, P. <i>et al.</i> (1989) The structural biology of CD2. <u>Immunol Rev. 111: 111</u><br>2. Beyers, A.D. <i>et al.</i> (1989) Activation of T lymphocytes via monoclonal antibodies<br>against rat cell surface antigens with particular reference to CD2 antigen. <u>Immunol</u><br><u>111: 59-77.</u>   | S                 |
| Storage                     | Store at +4°C.<br>DO NOT FREEZE.   |                   |

|                                  | This product should be stored undiluted.   |  |  |
|----------------------------------|--|--|--|
| Guarantee                        | 12 months from date of despatch  |  |  |
| Acknowledgements                 | This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts                                |  |  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #20471 available at:<br>https://www.bio-rad-antibodies.com/SDS/MCA1194SBUV605<br>20471 |  |  |
| Regulatory                       | For research purposes only   |  |  |

## **Related Products**

#### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

| North & South | Tel: +1 800 265 7376                 | Worldwide          | Tel: +44 (0)1865 852 700             | Europe            | Tel: +49 (0) 89 8090 95 21           |
|---------------|--------------------------------------|--------------------|--------------------------------------|-------------------|--------------------------------------|
| America       | Fax: +1 919 878 3751                 |                    | Fax: +44 (0)1865 852 739             |                   | Fax: +49 (0) 89 8090 95 50           |
|               | Email: antibody_sales_us@bio-rad.com |                    | Email: antibody_sales_uk@bio-rad.com |                   | Email: antibody_sales_de@bio-rad.com |
| To find a b   | atab (lat an a sifia alataab         | a at fau thia much | est interest und sum antime.         | a a such ta al at | , his rad antihadias com/datashasta  |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M410407:221028'

#### Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint