

Datasheet: MCA1195FT

| | |
|----------------------|----------------------------|
| Description: | MOUSE ANTI HUMAN CD21:FITC |
| Specificity: | CD21 |
| Other names: | CR2 |
| Format: | FITC |
| Product Type: | Monoclonal Antibody |
| Clone: | LB21 |
| Isotype: | IgG1 |
| Quantity: | 25 µg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | Neat - 1/10 |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Sheep, Mink, Bovine, Goat, Cat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
|-----------|-------------|---------------------|-------------------|
| | FITC | 490 | 525 |

Preparation

Purified IgG prepared by ion exchange chromatography from ascites

Buffer Solution

Phosphate buffered saline

| | |
|---------------------------------------|---|
| Preservative Stabilisers | 0.09% sodium azide (NaN ₃) 1% bovine serum albumin |
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml |
| Immunogen | Human IM9 cell line. |
| External Database Links | <p>UniProt: P20023 Related reagents</p> <p>Entrez Gene: 1380 CR2 Related reagents</p> |
| Synonyms | C3DR |
| RRID | AB_2085289 |
| Fusion Partners | Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/0 myeloma cell line. |
| Specificity | Mouse anti Human CD21 antibody, clone LB21 recognizes the human Complement receptor type 2, also known as CD21 or the Epstein-Barr virus receptor. CD21 is a ~14 kDa cell surface glycoprotein expressed by mature B cells and by follicular dendritic cells. The molecule acts as a receptor for complement components C3d, C3dg and iC3b, as well as for Epstein Barr Virus. It forms part of a large signal transduction complex in association with CD19. |
| Flow Cytometry | Use 10µl of the suggested working dilution to label 10 ⁶ cells or 100µl whole blood |
| References | <ol style="list-style-type: none"> 1. Furukawa, Y. <i>et al.</i> (2000) Frequent reversible membrane damage in peripheral blood B cells in human T cell lymphotropic virus type I (HTLV-I)-associated myelopathy/tropical spastic paraparesis (HAM/TSP). Clin Exp Immunol. 120 (2): 307-16. 2. Aichele, A. <i>et al.</i> (2006) Redox regulation of CD21 shedding involves signaling via PKC and indicates the formation of a juxtamembrane stalk. J Cell Sci. 119: 2892-902. 3. Sengstake, S. <i>et al.</i> (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. Int Immunol. 18: 1171-8. 4. Griebel, P.J. <i>et al.</i> (2007) Cross-reactivity of mAbs to human CD antigens with sheep leukocytes. Vet Immunol Immunopathol. 119: 115-22. 5. Aasted, B. <i>et al.</i> (2007) Reactivity of monoclonal antibodies to human CD antigens with cells from mink. Vet Immunol Immunopathol. 119: 27-37. 6. Sopp, P. <i>et al.</i> (2007) Cross-reactivity of mAbs to human CD antigens with cells from cattle. Vet Immunol Immunopathol. 119: 106-14. 7. Davis, W.C. <i>et al.</i> (2007) Use of flow cytometry to identify monoclonal antibodies that recognize conserved epitopes on orthologous leukocyte differentiation antigens in goats, llamas, and rabbits. Vet Immunol Immunopathol. 119: 123-30. 8. Angel, C.E. <i>et al.</i> (2009) Distinctive localization of antigen-presenting cells in human |

lymph nodes. [Blood. 113: 1257-67.](#)

9. Meister, R.K. *et al.* (2007) Progress in the discovery and definition of monoclonal antibodies for use in feline research. [Vet Immunol Immunopathol. 119: 38-46.](#)

10. Clutterbuck, E.A. *et al.* (2012) Pneumococcal conjugate and plain polysaccharide vaccines have divergent effects on antigen-specific B cells. [J Infect Dis. 205: 1408-16.](#)

11. Damgaard, B.M. *et al.* (2012) The effects of feed restriction on physical activity, body weight, physiology, haematology and immunology in female mink. [Res Vet Sci. 93: 936-42.](#)

12. Stachowiak, B. & Weingartl, H.M. (2012) Nipah virus infects specific subsets of porcine peripheral blood mononuclear cells. [PLoS One. 7 \(1\): e30855.](#)

13. Pooley, H.B. *et al.* (2022) Sheep vaccinated against paratuberculosis have increased levels of B cells infiltrating the intestinal tissue. [Vet Immunol Immunopathol. 252: 110482.](#)

| | |
|--------------------------------------|---|
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. |
| Guarantee | 12 months from date of despatch |
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1195FT 10041 |
| Regulatory | For research purposes only |

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M436997:250307'

Printed on 28 May 2025