

Datasheet: MCA1197B

Description:	MOUSE ANTI HUMAN CD7:Biotin
Specificity:	CD7
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	LT7
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Normal human peripheral blood lymphocytes.
External Database Links	UniProt: P09564 Related reagents

Entrez Gene:[924](#) CD7 [Related reagents](#)

RRID	AB_322287
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63.653 mouse myeloma cell line.
Specificity	Mouse anti Human CD7 antibody, clone LT7 recognizes the human T-cell antigen CD7, also known as TP41 or CD7. CD7 is a single chain cell surface glycoprotein of ~40 kDa, expressed by T cells, NK cells and thymocytes. CD7 is also expressed by tumor cells in some cases of acute myeloid leukaemia.
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none">Haynes, B.F. <i>et al.</i> (1989) Ontogeny of T-cell precursors: a model for the initial stages of human T-cell development. Immunol Today. 10 (3): 87-91.Aruffo, A. & Seed, B. (1987) Molecular cloning of two CD7 (T-cell leukemia antigen) cDNAs by a COS cell expression system. EMBO J. 6 (11): 3313-6.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1197B 10041
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)[HUMAN SEROBLOCK \(BUF070B\)](#)**North & South** Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: antibody_sales_us@bio-rad.com**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M429159:240320'

Printed on 20 Mar 2024

