

# Datasheet: MCA1209A700

Description:	MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 700		
Specificity:	MOUSE IgG1 NEGATIVE CONTROL		
Format:	ALEXA FLUOR® 700		
Product Type:	Negative/Isotype Control		
Isotype:	lgG1		
Quantity:	100 TESTS/1ml		

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
	Yes No Not Determined Suggested Dilution						
	Flow Cytometry	•			*		
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.						
Target Species	Negative Control						
Product Form	Purified IgG conjugated to Alexa Fluor® 700 - liquid						
Max Ex/Em	Fluorophore	Excitation M	lax (nm)	Emission Max (nm)			
	Alexa Fluor®700	702		723			
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% sodium azide (NaN <sub>3</sub> )						
Stabilisers	1% bovine serum albumin						
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml						
Immunogen	Human T lymphocytes.						

RRID	AB_844573				
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.				
Specificity	<b>Mouse IgG1 Negative Control antibody</b> is suitable for use as a negative control to assess non-specific binding of mouse IgG1 antibodies to target cells. Mouse IgG1 Negative Control antibody has been tested and found to be negative on the following rat cell types, peripheral blood leucocytes, thymocytes, splenocytes and macrophages.				
	Clone F8-11-13 recognizes the human CD45RA antigen, and therefore human leucocytes may be used as a positive control for this product. NOT SUITABLE FOR USE AS A NEGATIVE CONTROL ON HUMAN TISSUES				
Flow Cytometry	Use 10µl of the suggested working dilution to label $10^6$ cells in $100µ$ l				
References	<ol> <li>Weiss, D.J. <i>et al.</i> (2008) Bovine monocyte TLR2 receptors differentially regulate the intracellular fate of <i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> and <i>Mycobacterium avium</i> subsp. <i>avium</i>. J Leukoc Biol. 83 (1): 48-55.</li> <li>Chen, W. <i>et al.</i> (2009) Expression of toll-like receptor 4 in uvea-resident tissue macrophages during endotoxin-induced uveitis. Mol Vis. 15: 619-28.</li> <li>Safeukui I <i>et al.</i> (2015) Malaria induces anemia through CD8+ T cell-dependent parasite clearance and erythrocyte removal in the spleen. MBio. 6 (1) pii: e02493-14.</li> <li>Aricha, R. <i>et al.</i> (2016) Suppression of experimental autoimmune myasthenia gravis by autologous T regulatory cells. J Autoimmun. 67: 57-64.</li> <li>Wattegedera, S.R. <i>et al.</i> (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. Vet Res. 48 (1): 20.</li> <li>Stangl, H. <i>et al.</i> (2020) MHC/class-II-positive cells inhibit corticosterone of adrenal gland cells in experimental arthritis: a role for IL-1β, IL-18, and the inflammasome. Sci Rep. 10 (1): 17071.</li> <li>Terpeluk, R.E. <i>et al.</i> (2024) Supplementation of Foals with a Saccharomyces cerevisiae Fermentation Product Alters the Early Response to Vaccination <u>Animals. 14 (6): 960.</u></li> <li>Midavaine, É. <i>et al.</i> (2025) Humanized FccRI Expressed on Mouse Eosinophils Mediates IgE-Facilitated Eosinophil Antigen Presentation <u>Cells. 14 (4): 301.</u></li> </ol>				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in				
	frost-free freezers is not recommended. This product is photosensitive and should be protected from light.				
Guarantee	12 months from date of despatch				
Acknowledgements	This product is provided under an intellectual property licence from Life Technologies				

Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1209A700 10041
Regulatory	For research purposes only

### Related Products

#### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL: Alexa Fluor® 700 (MCA928A700)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	l.com	Email: antibody_sales_uk@bio-ra	id.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M408485:221012'

#### Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint