

Datasheet: MCA1209XZ

BATCH NUMBER 165947

Description:	MOUSE IgG1 NEGATIVE CONTROL:Preservative Free
Specificity:	MOUSE IgG1 NEGATIVE CONTROL
Format:	Preservative Free
Product Type:	Negative/Isotype Control
Isotype:	IgG1
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			*
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Negative Control
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human T lymphocytes.
RRID	AB_322266
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	<p>Mouse IgG1 Negative Control antibody is suitable for use as a negative control to assess non-specific binding of mouse IgG1 antibodies to target cells. Mouse IgG1 Negative Control antibody has been tested and found to be negative on the following rat cell types, peripheral blood leucocytes, thymocytes, splenocytes and macrophages.</p> <p>Clone F8-11-13 recognizes the human CD45RA antigen, and therefore human leucocytes may be used as a positive control for this product. NOT SUITABLE FOR USE AS A NEGATIVE CONTROL ON HUMAN TISSUES</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> 1. Weiss, D.J. <i>et al.</i> (2008) Bovine monocyte TLR2 receptors differentially regulate the intracellular fate of <i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> and <i>Mycobacterium avium</i> subsp. <i>avium</i>. J Leukoc Biol. 83 (1): 48-55. 2. Chen, W. <i>et al.</i> (2009) Expression of toll-like receptor 4 in uvea-resident tissue macrophages during endotoxin-induced uveitis. Mol Vis. 15: 619-28. 3. Safeukui I <i>et al.</i> (2015) Malaria induces anemia through CD8+ T cell-dependent parasite clearance and erythrocyte removal in the spleen. MBio. 6 (1) pii: e02493-14. 4. Aricha, R. <i>et al.</i> (2016) Suppression of experimental autoimmune myasthenia gravis by autologous T regulatory cells. J Autoimmun. 67: 57-64. 5. Wattedgedera, S.R. <i>et al.</i> (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. Vet Res. 48 (1): 20. 6. Stangl, H. <i>et al.</i> (2020) MHC/class-II-positive cells inhibit corticosterone of adrenal gland cells in experimental arthritis: a role for IL-1β, IL-18, and the inflammasome. Sci Rep. 10 (1): 17071. 7. Terpeluk, R.E. <i>et al.</i> (2024) Supplementation of Foals with a <i>Saccharomyces cerevisiae</i> Fermentation Product Alters the Early Response to Vaccination Animals. 14 (6): 960.
Storage	<p>Store at -20°C only.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA1209XZ

10162

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M408480:221012'

Printed on 25 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)