

# Datasheet: MCA1210F

Description:	MOUSE IgG2a NEGATIVE CONTROL:FITC
Specificity:	MOUSE IgG2a NEGATIVE CONTROL
Format:	FITC
Product Type:	Negative/Isotype Control
Isotype:	lgG2a
Quantity:	0.1 mg

### **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	•			*			
	Where this antibody has not been tested for use in a particular technique this does r necessarily exclude its use in such procedures.							
	* It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.							
Target Species	Negative Control							
Product Form	Purified IgG conjugate	d to Fluores	cein Isoth	niocyanate Isomer 1 (F	TTC) - liquid			
Max Ex/Em	Fluorophore	Excitation I	Max (nm)	Emission Max (nm)				
	FITC	490	)	525				
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant							
Buffer Solution	Phosphate buffered saline							
Preservative	0.09% sodium azide (NaN <sub>3</sub> )							
Stabilisers	1% bovine serum albumin							
Approx. Protein Concentrations	IgG concentration 0.1mg/ml							
Immunogen	Human T lymphocytes.							

RRID	AB_567343
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the NSI mouse myeloma cell line.
Specificity	<b>Mouse IgG2a Negative Control</b> is suitable for use as a matched isotype negative control reagent to assess the level of non-specific binding of mouse IgG2a monoclonal antibodies to rat cells. This antibody has been tested and found to be negative on the following rat cells – peripheral blood leucocytes, thymocytes, splenocytes and macrophages. Mouse IgG2a Negative Control ( <b>MCA1210</b> ) recognizes the human CD45 antigen and
	therefore human leucocytes may be used as a positive control for this product.
	THIS PRODUCT IS NOT SUITABLE FOR USE AS A NEGATIVE CONTROL ON HUMAN TISSUES.
Flow Cytometry	Use 10µl of the suggested working dilution to label $10^6$ cells in 100µl
References	<ol> <li>Fijak, M. <i>et al.</i> (2015) Influence of Testosterone on Inflammatory Response in Testicular Cells and Expression of Transcription Factor Foxp3 in T Cells. <u>Am J Reprod Immunol. 74</u> (<u>1): 12-25.</u></li> <li>Midavaine, É. <i>et al.</i> (2024) Discovery of a CCR2-targeting pepducin therapy for chronic pain. <u>Pharmacol Res. : 107242.</u></li> </ol>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1210F 10041
Regulatory	For research purposes only

# **Related Products**

### **Recommended Negative Controls**

#### MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

North & South	Tel: +1 800 265 7376 Wor	rldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com	ı	Email: antibody_sales_uk@bio-rad	l.com	Email: antibody_sales_de@bio-rad.com

#### Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint