

Datasheet: MCA1210XZ

Description:	MOUSE IgG2a NEGATIVE CONTROL:Preservative Free
Specificity:	MOUSE IgG2a NEGATIVE CONTROL
Format:	Preservative Free
Product Type:	Negative/Isotype Control
Isotype:	IgG2a
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Negative Control
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on protein A from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Human T lymphocytes.
RRID	AB_808774

Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the NSI mouse myeloma cell line.
Specificity	<p>Mouse IgG2a Negative Control is suitable for use as a matched isotype negative control reagent to assess the level of non-specific binding of mouse IgG2a monoclonal antibodies to rat cells. This antibody has been tested and found to be negative on the following rat cells – peripheral blood leucocytes, thymocytes, splenocytes and macrophages.</p> <p>Mouse IgG2a Negative Control (MCA1210) recognizes the human CD45 antigen and therefore human leucocytes may be used as a positive control for this product.</p> <p>THIS PRODUCT IS NOT SUITABLE FOR USE AS A NEGATIVE CONTROL ON HUMAN TISSUES.</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl.
References	<p>1. Fijak, M. <i>et al.</i> (2015) Influence of Testosterone on Inflammatory Response in Testicular Cells and Expression of Transcription Factor Foxp3 in T Cells. Am J Reprod Immunol. 74 (1): 12-25.</p> <p>2. Midavaine, É. <i>et al.</i> (2024) Discovery of a CCR2-targeting pepducin therapy for chronic pain. Pharmacol Res. : 107242.</p>
Storage	<p>Store at -20°C only.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA1210XZ 10162
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M408500:221012'

Printed on 23 May 2025