

Datasheet: MCA1212

BATCH NUMBER 149141

Description:	RAT IgG2a NEGATIVE CONTROL
Specificity:	RAT IgG2a NEGATIVE CONTROL
Format:	Purified
Product Type:	Negative/Isotype Control
Isotype:	IgG2a
Quantity:	1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*
Immunohistology - Frozen		■		
Immunohistology - Paraffin			■	
ELISA	■			
Immunoprecipitation			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. * It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.

Target Species	Negative Control
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml

Immunogen	Human lymphocytes.
RRID	AB_322674
Fusion Partners	Spleen cells from immunized DA rats were fused with cells of the rat Y3/Ag1.2.3. myeloma cell line.
Specificity	<p>Rat IgG2a Negative Control antibody is suitable for the assessment of the level of non-specific binding of rat IgG2a monoclonal antibodies to mouse cells.</p> <p>Test results indicate Rat IgG2a Negative Control antibody is also suitable for use as a negative control with canine cells.</p> <p>N.B. This antibody recognizes a human cell surface marker, and therefore is not suitable as a negative control in human cells or cell lines.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Sumagin, R. <i>et al.</i> (2008) Leukocyte-endothelial cell interactions are linked to vascular permeability via ICAM-1-mediated signaling. Am J Physiol Heart Circ Physiol. 295: H969-H977. 2. Chiu, W.C. <i>et al.</i> (2011) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. J Nutr Biochem. 20: 254-60. 3. Guilloteau, L.A. <i>et al.</i> (2003) Nramp1 is not a major determinant in the control of <i>Brucella melitensis</i> infection in mice. Infect Immun. 71: 621-8. 4. Stapleton, T.W. <i>et al.</i> (2000) Investigation of the regenerative capacity of an acellular porcine medial meniscus for tissue engineering applications. Tissue Eng Part A. 17: 231-42. 5. Park, S.W. <i>et al.</i> (2012) A1 adenosine receptor allosteric enhancer PD-81723 protects against renal ischemia-reperfusion injury. Am J Physiol Renal Physiol. 303: F721-32. 6. Schmidt, E.P. <i>et al.</i> (2012) The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. Nat Med. 18 (8): 1217-23. 7. McConnell, M.J. <i>et al.</i> (2009) H2-K(b) and H2-D(b) regulate cerebellar long-term depression and limit motor learning. Proc Natl Acad Sci U S A. 106: 6784-9. 8. Rabadi MM <i>et al.</i> (2016) Peptidyl arginine deiminase-4 deficient mice are protected against kidney and liver injury after renal ischemia and reperfusion. Am J Physiol Renal Physiol. Jun 22: ajprenal.00254.2016. [Epub ahead of print] 9. Rabadi, M.M. <i>et al.</i> (2019) Peptidyl arginine deiminase-4 exacerbates ischemic AKI by finding NEMO (NFkB Essential Modulator). Am J Physiol Renal Physiol. Apr 03 [Epub ahead of print]. 10. Han, S.J. <i>et al.</i> (2020) Renal proximal tubular NEMO plays a critical role in ischemic acute kidney injury. JCI Insight. 5 (19)Sep 17 [Epub ahead of print].
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p>

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1212 10041
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M364923:200529'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)