

## Datasheet: MCA1212

<b>Description:</b>	RAT IgG2a NEGATIVE CONTROL
<b>Specificity:</b>	RAT IgG2a NEGATIVE CONTROL
<b>Format:</b>	Purified
<b>Product Type:</b>	Negative/Isotype Control
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	1 ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*
Immunohistology - Frozen		■		
Immunohistology - Paraffin			■	
ELISA	■			
Immunoprecipitation			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. \* It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.

<b>Target Species</b>	Negative Control
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative</b>	0.09% Sodium Azide
<b>Stabilisers</b>	1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Human lymphocytes.
<b>RRID</b>	AB_322674
<b>Fusion Partners</b>	Spleen cells from immunized DA rats were fused with cells of the rat Y3/Ag1.2.3. myeloma cell line.
<b>Specificity</b>	<b>Rat IgG2a Negative Control antibody</b> is suitable for the assessment of the level of non-specific

binding of rat IgG2a monoclonal antibodies to mouse cells.

Test results indicate Rat IgG2a Negative Control antibody is also suitable for use as a negative control with dog cells.

**N.B.** This antibody recognizes a human cell surface marker, and therefore is not suitable as a negative control in human cells or cell lines.

---

**Flow Cytometry** Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

---

**References**

1. Sumagin, R. *et al.* (2008) Leukocyte-endothelial cell interactions are linked to vascular permeability via ICAM-1-mediated signaling. [Am J Physiol Heart Circ Physiol. 295: H969-H977.](#)
2. Chiu, W.C. *et al.* (2011) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. [J Nutr Biochem. 20: 254-60.](#)
3. Guilloteau, L.A. *et al.* (2003) Nramp1 is not a major determinant in the control of *Brucella melitensis* infection in mice. [Infect Immun. 71: 621-8.](#)
4. Stapleton, T.W. *et al.* (2000) Investigation of the regenerative capacity of an acellular porcine medial meniscus for tissue engineering applications. [Tissue Eng Part A. 17: 231-42.](#)
5. Park, S.W. *et al.* (2012) A1 adenosine receptor allosteric enhancer PD-81723 protects against renal ischemia-reperfusion injury. [Am J Physiol Renal Physiol. 303: F721-32.](#)
6. Schmidt, E.P. *et al.* (2012) The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. [Nat Med. 18 \(8\): 1217-23.](#)
7. McConnell, M.J. *et al.* (2009) H2-K(b) and H2-D(b) regulate cerebellar long-term depression and limit motor learning. [Proc Natl Acad Sci U S A. 106: 6784-9.](#)
8. Rabadi MM *et al.* (2016) Peptidyl arginine deiminase-4 deficient mice are protected against kidney and liver injury after renal ischemia and reperfusion. [Am J Physiol Renal Physiol. Jun 22: ajrenal.00254.2016. \[Epub ahead of print\]](#)
9. Rabadi, M.M. *et al.* (2019) Peptidyl arginine deiminase-4 exacerbates ischemic AKI by finding NEMO (NFκB Essential Modulator). [Am J Physiol Renal Physiol. Apr 03 \[Epub ahead of print\].](#)

---

**Storage** Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

---

**Regulatory** For research purposes only

---

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M364923:200529'

Printed on 11 Aug 2020

