

Datasheet: MCA1212A405

BATCH NUMBER 1710

Description:	RAT IgG2a NEGATIVE CONTROL:Alexa Fluor® 405
Specificity:	RAT IgG2a NEGATIVE CONTROL
Format:	ALEXA FLUOR® 405
Product Type:	Negative/Isotype Control
Isotype:	lgG2a
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. * It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody

Target Species	Negative Control						
Product Form	Purified IgG conjuga	Purified IgG conjugated to Alexa Fluor® 405 - liquid					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	Alexa Fluor®405	401	421				
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered	Phosphate buffered saline					
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum	n Albumin					
Approx. Protein Concentrations	IgG concentration 0.0	05 mg/ml					
Immunogen	Human lymphocytes						

RRID AB 567360 **Fusion Partners** Spleen cells from immunized DA rats were fused with cells of the rat Y3/Ag1.2.3. myeloma cell line. **Specificity** Rat IgG2a Negative Control antibody is suitable for the assessment of the level of non-specific binding of rat IgG2a monoclonal antibodies to mouse cells. Test results indicate Rat IgG2a Negative Control antibody is also suitable for use as a negative control with canine cells. N.B. This antibody recognizes a human cell surface marker, and therefore is not suitable as a negative control in human cells or cell lines. References 1. Sumagin, R. et al. (2008) Leukocyte-endothelial cell interactions are linked to vascular permeability via ICAM-1-mediated signaling. Am J Physiol Heart Circ Physiol. 295: H969-H977. 2. Chiu, W.C. et al. (2011) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. J Nutr Biochem. 20: 254-60. 3. Guilloteau, L.A. et al. (2003) Nramp1 is not a major determinant in the control of Brucella melitensis infection in mice. Infect Immun. 71: 621-8. 4. Stapleton, T.W. et al. (2000) Investigation of the regenerative capacity of an acellular porcine medial meniscus for tissue engineering applications. Tissue Eng Part A. 17: 231-42. 5. Park, S.W. et al. (2012) A1 adenosine receptor allosteric enhancer PD-81723 protects against renal ischemia-reperfusion injury. Am J Physiol Renal Physiol. 303: F721-32.

- 6. Schmidt, E.P. et al. (2012) The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. Nat Med. 18 (8): 1217-23.
- 7. McConnell, M.J. et al. (2009) H2-K(b) and H2-D(b) regulate cerebellar long-term depression and limit motor learning. Proc Natl Acad Sci U S A. 106: 6784-9.
- 8. Rabadi MM et al. (2016) Peptidyl arginine deiminase-4 deficient mice are protected against kidney and liver injury after renal ischemia and reperfusion. Am J Physiol Renal Physiol. Jun 22: ajprenal.00254.2016. [Epub ahead of print]
- 9. Rabadi, M.M. et al. (2019) Peptidyl arginine deiminase-4 exacerbates ischemic AKI by finding NEMO (NFkB Essential Modulator). Am J Physiol Renal Physiol. Apr 03 [Epub ahead of print].
- 10. Han, S.J. et al. (2020) Renal proximal tubular NEMO plays a critical role in ischemic acute kidney injury. JCI Insight. 5 (19)Sep 17 [Epub ahead of print].

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use

12 months from date of despatch
The Alexa Fluor® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays, and are covered by pending and issued patents
Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1212A405 10041
For research purposes only

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide Fax: +44 (0)1865 852 739

Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M364924:200529'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint