

Datasheet: MCA1212F

Description:	RAT IgG2a NEGATIVE CONTROL:FITC
Specificity:	RAT IgG2a NEGATIVE CONTROL
Format:	FITC
Product Type:	Negative/Isotype Control
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10*

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Negative Control		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Human lymphocytes.		

Fusion Partners Spleen cells from immunized DA rats were fused with cells of the rat Y3/Ag1.2.3. myeloma cell line.

Specificity **Rat IgG2a Negative Control antibody** is suitable for the assessment of the level of non-specific binding of rat IgG2a monoclonal antibodies to mouse cells.

Test results indicate Rat IgG2a Negative Control antibody is also suitable for use as a negative control with canine cells.

N.B. This antibody recognizes a human cell surface marker, and therefore is not suitable as a negative control in human cells or cell lines.

References

1. Stapleton, T.W. *et al.* (2000) Investigation of the regenerative capacity of an acellular porcine medial meniscus for tissue engineering applications. [Tissue Eng Part A. 17: 231-42.](#)
 2. Guilloteau, L.A. *et al.* (2003) Nramp1 is not a major determinant in the control of *Brucella melitensis* infection in mice. [Infect Immun. 71: 621-8.](#)
 3. Sumagin, R. *et al.* (2008) Leukocyte-endothelial cell interactions are linked to vascular permeability via ICAM-1-mediated signaling. [Am J Physiol Heart Circ Physiol. 295: H969-H977.](#)
 4. McConnell, M.J. *et al.* (2009) H2-K(b) and H2-D(b) regulate cerebellar long-term depression and limit motor learning. [Proc Natl Acad Sci U S A. 106: 6784-9.](#)
 5. Chiu, W.C. *et al.* (2011) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. [J Nutr Biochem. 20: 254-60.](#)
 6. Schmidt, E.P. *et al.* (2012) The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. [Nat Med. 18 \(8\): 1217-23.](#)
 7. Park, S.W. *et al.* (2012) A1 adenosine receptor allosteric enhancer PD-81723 protects against renal ischemia-reperfusion injury. [Am J Physiol Renal Physiol. 303: F721-32.](#)
 8. Rabadi, M. *et al.* (2016) Peptidyl arginine deiminase-4-deficient mice are protected against kidney and liver injury after renal ischemia and reperfusion. [Am J Physiol Renal Physiol. 311 \(2\): F437-49.](#)
 9. Rabadi, M.M. *et al.* (2019) Peptidyl arginine deiminase-4 exacerbates ischemic AKI by finding NEMO. [Am J Physiol Renal Physiol. 316 \(6\): F1180-F1190.](#)
 10. Han, S.J. *et al.* (2020) Renal proximal tubular NEMO plays a critical role in ischemic acute kidney injury. [JCI Insight. 5 \(19\): e139246.](#)
 11. Ono, Y. *et al.* (2018) CD11c+ M1-like macrophages (M ϕ s) but not CD206+ M2-like M ϕ are involved in folliculogenesis in mice ovary. [Sci Rep. 8 \(1\): 8171.](#)
-

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1212F 10041
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA6005F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M408534:221013'

Printed on 19 Jan 2024